



Not Just Cutting It, a Successful Interprofessional Cooperation for the Treatment of Intractable Cancer Pain

Uri Hochberg^{1,2*}, Rotem Tellem^{1,3}, Ido Strauss^{1,4,5}

¹Department of Medicine, Tel Aviv University, Israel

²Department of Pain Medicine, Tel Aviv University, Israel

³Department of Palliative Care, Tel Aviv University, Israel

⁴Department of Neurosurgery Unit, Tel Aviv University, Israel

⁵Department of Neurosurgery, Tel Aviv University, Israel

Short Communication

The modern era in oncology brought a major change in the composition of patients treated for cancer in the oncology divisions. New cutting-edge anticancer biological and immunological treatments provide long term survival even for advanced stage 4 cancer patients.

One of the major challenges in the care of advanced cancer patients is pain management. These patients often suffer from severe pain that is refractory to standard medical treatment according to the WHO guidelines and require out of the box solutions. The presence of pain influences greatly their quality of life and functional status.

The consideration of interventional pain procedures as a "fourth" step to the WHO ladder has been widely discussed in the literature [1]. The most commonly applied interventions are intrathecal or epidural drug delivery systems and/or peripheral nerve blocks and ablations. However, these neuromodulatory techniques although non-destructive and reversible, require ongoing maintenance, and also carry specific risks such as infection, especially in immunocompromised and cachectic patients. Patients, who suffer from refractory cancer pain, may benefit from targeted neurosurgical procedures that selectively intervene along the central pain pathways. The neurosurgical ablative armamentarium to treat pain includes a variety of procedures that can be tailored to the patient's specific pain syndrome. Patients with localized unilateral pain can be treated by disconnection of the spinothalamic pain pathways using percutaneous cordotomy or stereotactic mesencephalotomy. Patients who suffer from diffuse pain can undergo stereotactic cingulotomy that decreases the overall suffering associated with the perception of pain. These procedures are minimally invasive, can be performed without general anesthesia and most importantly achieve immediate and long lasting pain relief.

Appropriate patient selection is essential for the success of these procedures and is not a one person's job. In the last three years we have incorporated the use of these interventions in our oncology division. We have formed an interdisciplinary service comprised of a palliative care physician, a neurosurgeon and a pain specialist. The service evaluates advanced cancer patients who suffer from intractable pain, referred from their oncology or palliative care providers. Our patient selection process includes a thorough pain assessment, revision of the different modalities already attempted to alleviate pain and an honest and detailed discussion about the goals is done with the patients and their families. After this evaluation, the team is able to offer a neurosurgical intervention that is most appropriate to the patients' medical status, their goals of care and in accordance to their oncological treatment plan.

In the course of our work we have managed to improve the pain and the quality of life of some of the most complex and advanced cancer patients, previously diagnosed with intractable pain. In many patients we have seen a marked reduction in opioids analgesics doses, improved function and even the ability to participate in oncology treatments that were previously intolerable due to pain [2,3]. With shared, wholistic, decisions making process, we have managed to offer the most complex patients a unique, "out of the box", solution and exemplary palliation in accordance to the ethos of modern palliative care profession: "You matter because you are you, and you matter to the end of your life. We will do all we can not only to help you die peacefully, but also to live until you die." - Dame Cicely Saunders, founder of the modern hospice movement.

OPEN ACCESS

*Correspondence:

Uri Hochberg, Department of Medicine,
Tel Aviv University, Israel,
E-mail: urihochberg@hotmail.com

Received Date: 14 Aug 2018

Accepted Date: 03 Sep 2018

Published Date: 12 Sep 2018

Citation:

Hochberg U, Tellem R, Strauss I.
Not Just Cutting It, a Successful
Interprofessional Cooperation for the
Treatment of Intractable Cancer Pain.
Clin Oncol. 2018; 3: 1519.

Copyright © 2018 Uri Hochberg. This
is an open access article distributed
under the Creative Commons Attribution
License, which permits unrestricted
use, distribution, and reproduction in
any medium, provided the original work
is properly cited.

References

1. Vissers KC, Besse K, Wagemans M, Zuurmond W, Giezeman MJ, Lataster A, et al. Pain in patients with cancer. *Pain Pract*. 2011;11(5):453-75.
2. Strauss I, Berger A, Ben Moshe S, Arad M, Hochberg U, Gonen T, et al. Double Anterior Stereotactic Cingulotomy for Intractable Oncological Pain. *Stereotact Funct Neurosurg*. 2017;95(6):400-8.
3. Strauss I, Berger A, Arad M, Hochberg U, Tellem R. O-Arm-Guided Percutaneous Radiofrequency Cordotomy. *Stereotact Funct Neurosurg*. 2017;95(6):409-16.