



Innovative Exploratory Clinical Approaches for Relapsed and/or Refractory Metastatic Ewing's Sarcoma

Ghisoli M^{1,2,3}, Manning L⁴, Senzer N^{1,4,5} and Nemunaitis J^{1,2,3,4,5*}

¹Mary Crowley Cancer Research Centers, USA

²Medical Oncology and Hematology, Texas Oncology, USA

³Medical City Dallas Hospital, USA

⁴Gradalis, Inc., USA

⁵Strike Bio, USA

Abstract

Relapsed and/or refractory Ewing's Sarcoma is a devastating pediatric disease with rapid progression and oftentimes severe side effects related to generally ineffective high dose multi-agent chemotherapy. Eighty five percent of diagnosed Ewing's Sarcoma is characterized by EWS/FLI1 fusion gene expression that provides a unique opportunity for targeted therapeutics development. The EWS/FLI1 gene is a "driver gene" with transformative potential and integral to Ewing's cancer progression. Although encoding a transcription factor, which is pharmacologically "undruggable", it connects with potentially targetable molecular signals and, in addition, as a fusion gene along with accompanying tumor specific mutations provides unique neoantigens some of which process into immunogenic epitope. Very few cutting edge advances for the management and control of Ewing's Sarcoma have been made in the last 20 years due, in part, to low incidence (one case per million people), a narrow therapeutic window, and a limited availability of tissue suitable for biomarker studies. However, recent advances in DNA/RNA manipulation [CRISPR and RNA interference (siRNA)] as well as in molecular and immune technologies have transformed both the understanding of signaling pathways and molecular mechanisms of actions and, consequently, the approach to target identification. We review the innovative exploratory approaches to five unique therapies (a EWS-FLI1 co-activator, the EWS-FLI1 fusion gene itself, a signaling receptor, a DNA damage repair component, and the antigenic matrix) currently undergoing clinical assessment in Ewing's Sarcoma for which preliminary preclinical and clinical results suggest therapeutic benefit.

OPEN ACCESS

*Correspondence:

John Nemunaitis, Medical Oncology and Hematology, Mary Crowley Cancer Research Centers, 12222 Merit Drive Suite 1500, Dallas, Texas 75251, USA, Tel: 214-658-1964; Fax: 214-658-1992; E-mail: jnemunaitis@marycrowley.org

Received Date: 22 Jul 2016

Accepted Date: 17 Aug 2016

Published Date: 08 Sep 2016

Citation:

Ghisoli M, Manning L, Senzer N, Nemunaitis J. Innovative Exploratory Clinical Approaches for Relapsed and/or Refractory Metastatic Ewing's Sarcoma. *Clin Oncol.* 2016; 1: 1079.

Copyright © 2016 Nemunaitis J. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Background

Ewing's Family Sarcoma is a highly aggressive and malignant bone tumor that metastasizes frequently. The median age of diagnosis is 14-15 years [1,2] and the incidence rate is 1 case per million people in the United States but as high as 9-10 cases per million in the 10 to 19 year old age range [3].

Eighty five percent of Ewing's Sarcoma patients show a balanced translocation of the EWS gene at chromosome 22q12 with the FLI1 gene at chromosome 11q24 [4]. The EWS-FLI1 translocation can occur at one of several different gene fusion breakpoint sites. Most frequently seen are Type 1 (accounting for 60%) and Type 2 (25%). In Type 1 EWS-FLI1, exons 1-7 of EWS are fused with exons 6-9 of the FLI1 gene [5]. Some of the early clinical studies suggested a relationship of fusion type to rapid progression of disease [6,7] that more recent studies have not confirmed [8]. The Type 2 fusion comprised of EWS exons 1-7 juxtaposed to exons 5-9 of FLI1 [9], is associated with a higher Ewing's Sarcoma proliferation rate that may or may not have clinical significance. Ten to 15 percent of patients with Ewing's Sarcoma show other translocations; the EWS-ERG gene fusion (t(22;21)(q22;q12)) [10-12], and the less frequent EWS-ETS fusion group (EWS-ETV1 t(7;22), EWS-ETV4 t(17;22), EWS-FEV t(2;22)) [13-16]. Methodologies used to categorize the EWS-FLI1 translocations include real-time polymerase chain reaction (RT-PCR), fluorescence in situ hybridization (FISH), and next generation sequencing (NGS) [17,18].

At diagnosis, less than 25% of patients present with metastatic disease, however up to 90% of Ewing's adolescents eventually experience either disease progression or relapse after frontline treatment [3,19]. The most important prognostic factor for survival following failure of first-line

Table 1: Treatment options for patients with advanced, refractory or recurrent Ewing's Sarcoma. Comparison of objective response rate (ORR) and progression free survival (PFS).

Treatment	Mechanism	No. of EWS Patients (n)	ORR	PFS	OS	Reference
Ganitumumab (AMG479)	IGFR-1R Inhibitor	22	6%	7.9 mo		(Tap, Demetri et al. 2012)
R1507	IGFR-1R Inhibitor	109	10%	-		(Pappo, Patel et al. 2011)
Figitumumab (CP-751,871)	IGFR-1R Inhibitor	16	13%	-		(Olmos, Postel-Vinay et al. 2010)
Figitumumab (CP-751,871)	IGFR-1R Inhibitor	106	14%	1.9 mo		(Juergens, Daw et al. 2011)
Linsitinib	IGFR-1R Inhibitor	ongoing	-	-		EUROSARC trial
Olaparib	PARP-Inhibitor	12	0%	1.5 mo		(Choy, Butrynski et al. 2014)
Niraparib/Temozolomide	PARP-Inhibitor/alkylating agent	ongoing	-	-		(Wilcoxon, Brooks et al. 2013)
Olaparib/Temozolomide	PARP-Inhibitor/alkylating agent	ongoing	-	-		(Engert, Schneider et al. 2015)
YK-4-279	EWS-FLI1 protein RHA-binding inhibitor	ongoing	-	-		(Erkizan, Kong et al. 2009)
pbi-shRNA EWS/FLI1 Lipoplex (LPX)	bi-functional sh-RNA targeting EWS/FLI1 fusion protein	ongoing	-	-		(Rao, Jay et al. 2016)
Vigil	GMCSF/bi-shRNA ^{luc} DNA constructed autologous tumor vaccine	16	6%	-	24 mo	(Ghisoli, Barve et al. 2016)

treatment is relapse-time ≤ 2 years, which is associated with 5-year event free survival (EFS) of only 5% [2,20]. After second-line treatment, only 9-13% of patients will achieve second-line remission and most of these relapse rapidly on completion [20-22]. In a retrospective analysis of 195 advanced, \geq second-line therapy, metastatic Ewing's Sarcoma patients 86% did not achieve second remission and of those (n=26) who did so, the majority either re-relapsed or died within the year [20]. There are no standard of care (SOC) NCI recommendations for second-line treatment with advanced Ewing's Sarcoma, although multi-agent regimens including irinotecan, temozolomide, topotecan, and cyclophosphamide are commonly utilized today. In addition to relative ineffectiveness, the use of intensive chemotherapy in both frontline and second-line treatments of Ewing's Sarcoma is associated with a severe toxicity and morbidity.

Given the limitation of cumulative toxicity associated with chemotherapy and the emergence of resistance, it is not surprising that third-line management is even more challenging. Single or combined chemotherapy regimens only show limited response (in both rate and durability). Some regimens include high dose ifosfamide or gemcitabine/docetaxel [23-25]. Unfortunately, to date there has been no significant survival advantage to any \geq third-line therapy for patients with relapsed or refractory, disease Ewing's Sarcoma [26].

Bottom line: there is a need for both innovative treatment approaches and a greater array of therapeutic options in second- and third-line management of Ewing's Sarcoma. In the following we focus on experimental therapies currently in clinical trial for \geq third-line treatment of metastatic refractory and/or relapsed Ewing's Sarcoma (Table 1).

YK-4-279

YK-4-279 [27] is a small molecule that interacts with RNA Helicase A (RHA, encoded by the DHX9 gene) thereby affecting EWS/FLI1 signaling activity. The EWS/FLI1 fusion protein binds RHA in a unique region not targeted by other transcriptional proteins and thereby inhibits helicase activity in a dose dependent manner [28]. YK-4-279 binds to RHA adjacent to its helicase domain and to an as yet not completely specified region on the EWS/FLI1 fusion protein to disinhibit helicase activity but without affecting ATPase activity. Erkizan and colleagues have shown that YK-4-279 may significantly shift the RNA binding profiles of both EWS/FLI1 and RHA. RHA is a

transcriptional co-activator regulating both transcription and mRNA splicing and plays a role in both oncogenesis and tumor maintenance. Whatever the dominant mechanism, YK-4-279 results in inhibition of oncogenic activity and activation of caspase-3-induced apoptosis *in vitro* in a range of Ewing's Sarcoma cell lines *in vitro* (TC32, A4573, TC71, and ES925 in Figure 1) and *in vivo* [29-31]. Preclinical data suggest a chimeric structure of the small molecule with (S) and (R)-enantiomers with a (S)-YK-4-279 enantiomer-specific effect in EWS/FLI1 cells [29,30], (Figure 1). Disruption of protein-protein interactions, such as the transcription complex in Ewing's Sarcoma cells, comprising RNA polymerase II, CREB-binding protein (CBP), and RNA Helicase a (RHA) [32], thus seems to be a reasonable goal for therapeutic effectiveness. Interestingly, the same small molecule also demonstrates activity in other tumors with ETS family translocations such as ETV1 fusion-positive prostate cancer. The preclinical data demonstrates YK-4-279 inhibition of tumor growth as well as decreased motility and invasion of prostate cancer xenografts [33]. While EWS/FLI1 Types 1 and 2 are the two most common translocations in Ewing's Sarcoma and a further 10% of the patients





Figure 1: YK-4-279 reverses the inhibitory effect of EWS-FLI1 on RHA in an enantiomer-specific manner. Immobilized fulllength RHA on a CM5 chip and purified EWS-FLI1 were used in RHA assays. Figure shows recombinant RHA activity and ssRNA product in presence of either single recombinant RHA, or recombinant RHA / recombinant EWS-FLI1/YK-4-279, or recombinant RHA/recombinant EWS-FLI1/S-YK-4-279, or recombinant RHA/recombinant EWS-FLI1/R-YK-4-279. Results were plotted over time; x-axis represents time (sec) and y-axis is percent (product in helicase assay). Both racemic YK-4-279 and (S)-YK-4-279 disinhibited the helicase reaction, showing restoration of 80% helicase activity, while (R)-YK-4-279 does not restore RHA activity (Erkizan, Schneider et al. 2015).




reveal a EWS/ERG gene fusion, there is a less frequent population that presents with EWS/ETS-like fusions [13]. A similarly modified small molecular inhibitor, TK216, is listed on clinicaltrials.gov as a Phase I trial opportunity in patients with advanced Ewing's Sarcoma.

Bi-sh (Bifunctional Short Hairpin) RNA EWS-FLI1 Type 1 Lipoplex (LPX)

The dual stem-loop bi-shRNA EWS-FLI1 (Type 1) incorporated into the pUMVC3 plasmid construct transcribes both siRNA and miRNA-like effectors that target the identical junction region of the EWS-FLI1 fusion gene encoded mRNA [34-36]. The plasmid is systemically delivered in a DOTAP (cationic lipid dioleoyl trimethyl ammonium propane)/cholesterol delivery vehicle [37] as a lipoplex (LPX). This RNAi technology obviates the inherent difficulty of targeting the undruggable EWS-FLI1 protein and by targeting the Type 1 breakpoint to down regulates the expression of the EWS/FLI1 encoded mRNA and protein. Preclinical testing *in vitro* and *in vivo* demonstrated 85-92% type-specific knockdown of target protein [38]. Bi-shRNA simultaneously induces RISC (RNA induced silencing complex)-cleavage-dependent mRNA degradation and RISC-cleavage-independent degradation of same nucleotide sequence. Bi-sh RNA affects targeted protein down regulation at a 5-log lower dose in comparison to si-RNA targeting the same strand sequence [34]. Furthermore, the bi-sh RNA EWS/FLI1 Type 1 dual effect or target sequence-specific activity limits the potential for off-target effects against "non" Type 1 Ewing's Sarcoma fusion constructs [38]. A significant tumor growth delay and survival advantage *in vivo* was demonstrated in human Type 1 EWS/FLI1 cells treated with the bi-sh RNA EWS/FLI1 Type 1 LPX (Figure 2). As hypothesized, the specificity of fusion gene encoded protein knockdown as compared with wild type FLI1 protein which was not knocked down was confirmed in HEK 293 cells that contained both the wild type EWS and wild type FLI1. Figure 3 shows the predicted specificity by comparing the response of SK-N-MC cells containing the EWS/FLI1 Type 1 fusion gene to the response of HEK-293 cells with wild types EWS and FLI1 genes. Importantly, GMP (good manufacturing practice) safety testing in large animals revealed excellent tolerability [38] at active human equivalent dose ranges. Clinical investigation has been initiated in refractory/relapsed Ewing's Sarcoma patients with the Type 1 EWS/FLI1 gene fusion (ClinicalTrials.gov).

IGF-1R Inhibitors

The EWS/FLI1 translocation is associated with dysregulation of the insulin growth factor receptor (IGF-1R) pathway. An oncogenic role for co-activation of IGF-1R signaling has been suggested based on preclinical assessment [39]. In Ewing's Sarcoma there is evidence for autocrine activation of the IGF-1R pathway as well as EWS-FLI1 induced over expression of the caveolin-1 membrane transport protein by way of which IGF-1R internalizes [40]. In addition, the fusion product represses IGF1BP3 that binds IGF-1 in the plasma thereby up regulating ligand-receptor induced signaling. Enhanced IGF-1R mediated activity can stream through two parallel pathways: 1) the PI3k/AKT pathway inhibiting apoptosis, increasing protein synthesis and promoting glucose metabolism and 2) the Ras/MAPK pathway promoting cancer cell proliferation [41]. Preclinical studies in cancer have demonstrated the relationship of the intrinsic tyrosine kinase activity of the IGF-1R with tumor proliferative and anti-apoptotic activity. Therefore, based on rationale and preclinical support, trials of therapeutic anti-tumor targeting of IGF-1R have been initiated [42-44]. In fact, IGF-1R inhibitor therapy has demonstrated benefit in



subsets of patients with Ewing's Sarcoma [45,46]. However, activity is limited with response rates between 6% to 14% in Phase I and II clinical trials of patients with advanced Ewing's Sarcoma who are generally undergoing second or third-line treatment (Table 1) [47-50]. Lack of effectiveness can be ascribed to multiple mechanisms including, but not limited to, lower than expected surface membrane receptor density, release from negative feedback inhibition pathways, and alternative signaling via enhanced insulin receptor (IR)-A homodimer formation.

Tap et al. [50] enrolled 38 patients including 22 with Ewing's Sarcoma and 16 with desmoplastic small round cell tumor (DSRCT) into a Phase II study to test the monoclonal antibody (MAb) IGF-1R inhibitor ganitumab (AMG479). They observed an objective response rate of 6%. Forty-nine percent of the patients had stable disease (SD), only four of whom achieved SD for ≥ 24 weeks. In another trial 115 patients with refractory or recurrent Ewing's Sarcoma were treated with a different MAb IGF-1R inhibitor, R1507, achieving an overall response of 10% with a median OS of 7.6 months (95% CI, 6 to 9.7 months) [49].

A third MAb IGF-1R inhibitor figitumumab was tested in a 29 patient study (16 of whom had Ewing's Sarcoma). Two patients (12.5%), both with Ewing's Sarcoma, had partial responses and

37.5% (6/16) stable disease [47]. A Phase I/II study was subsequently conducted to investigate the safety and effectiveness of figitumumab in patients with Ewing's Sarcoma [48]. However, only 1 of 31 patients achieved partial response (PR). Despite limited efficacy a Phase II study was performed involving 106 heavily pre-treated ($\geq 1 - 4$ prior lines of chemotherapy) patients with refractory or recurrent Ewing's Sarcoma. Fourteen percent (15/106) of these patients achieved PR and 23% (25/106) stable disease (SD); the median progression free survival was 1.9 months, and median overall survival 8.9 months (95% CI, 7.2 to 11.1) [48]. Although treatment was generally well tolerated, three cases of leukemia were observed; one attributed to figitumumab (after one cycle in a patient previously treated with doxorubicin and etoposide), another to concurrent rapamycin, and a third to prior etoposide. Figitumumab associated leukemia has not been reported in studies of other cancer types with IGR-1R inhibitors. Even though therapy related myelodysplasia and acute myeloid leukemia are known adverse events in Ewing's Sarcoma patients treated with chemotherapy, particularly in association with ifosfamide [51-53], the possibility of figitumumab associated leukemia cannot be entirely excluded.

These trials suggest mild to moderate clinical activity in Ewing's Sarcoma and establish a reasonable safety profile. Few of the studies incorporated predictive biomarkers (e.g., increased expression of IRS2 (insulin receptor substrate), IR, growth hormone (GH) and decreased expression of IGF-binding protein-5) to help identify those patients with a higher likelihood of response. Based on pathway analysis, rationale based clinical trials involving IGF-1R inhibitors are being explored; the concurrent administration of IGF-1R and mTOR inhibitors to attenuate negative feedback inhibition and dual IGF-1R/IR kinase inhibitors (e.g., Linsitinib) to block compensatory increased expression of IR [54], (ClinicalTrials.gov ID: NCT02546544).

PARP Inhibitors

Poly-ADP-ribose-polymerases (PARP1 and PARP2) are comprised of enzymes that transfer ADP-ribose onto target proteins (PARylation), thereby modifying a wide range of cellular processes including genome maintenance, transcriptional regulation, cell cycle control, proliferation, differentiation, necrosis and apoptosis [55,56]. PARP1, activated by DNA damage, binds to DNA single-strand breaks (SSB) and double strand breaks (DSB), then catalyzes and promotes multiple DNA repair processes [56]. Patients with cancer related mutations in BRCA1 or BRCA2, suppressor proteins involved in DSB repair, demonstrate enhanced sensitivity to PARP1-inhibitors with a consequent increase in apoptosis [57]. The anti-tumor activity of PARP-inhibitors has been confirmed in BRCA-mutant breast, ovary and prostate cancers [58-60]. Garnett et al. [61] was able to show Ewing's Sarcoma cell line sensitivity to PARP1-inhibitors by way of decreased viability of EWS/FLI1 cancer cells (Figure 4). Brenner and colleagues [62] hypothesized a reciprocal positive feedback loop in Ewing's Sarcoma; EWS-FLI1 protein driving PARP1 expression, the latter then facilitating EWS-FLI1 transcriptional activation. In addition, 7% of Ewing's Sarcoma patients have been shown to have BRCA2 mutations [63].

Choy et al. [64] conducted a two-part Phase II clinical trial of olaparib enrolling 12 patients with advanced Ewing's Sarcoma progressing following chemotherapy. None of the patients had an objective response (RECIST 1.1 criteria PR/CR). Four of the 12 patient's sustained SD for 10.9 to 17.9 weeks. Median time to progression was 5.7 weeks [64]. Based on the results of Part 1,




Figure 4: Sensitivity to olaparib of EWS-FLI1 (and FUS-CHOP) transformed mouse mesenchymal cells compared to the SK-N-MC cell line (which harbors the EWS-FLI1 fusion (Garnett, Edelman et al. 2012).




Figure 5: Temozolomide enhanced PARP-trapping in Ewing Sarcoma cell lines. Fold-induction of caspase 3/7 activation in EWS-cells following treatment with vehicle, olaparib single, temozolomide single, olaparib/temozolomide combination (Gill, Travers et al. 2015).

enrollment to Part 2 was put on hold. However, reanalysis and future assessment of PARP1 inhibitor effectiveness as well as protocol design need to take into account the mechanistic differences between two recently described classes of PARP1 inhibitors; 1) those that effect catalytic inhibition of PARP enzyme activity and 2) those that result in formation of PARP-traps that function as cytotoxic PARP-DNA complexes [65]. On the basis of catalytic inhibitory activity, the effectiveness of the three clinical PARP inhibitors ranks as follows: olaparib>veliparib>niraparib. Based on active cytotoxic PARP-DNA formation the ranking is niraparib>olaparib>veliparib.

Drug-sensitivity testing of PARP inhibition in combination with various S-phases DNA damaging agents in Ewing’s Sarcoma cell lines [66] showed enhanced activity with the combination of olaparib and temozolomide. Engert et al. [67] demonstrated that combined olaparib and temozolomide up-regulated the pro-apoptotic proteins BAX and BAK and caspase activation. Synergistic activity was also demonstrated with the combination of niraparib and temozolomide [68]. These results are presumably due to interaction with the normally sublethal effects of temozolomide induced lesions insofar as Ewing’s Sarcoma cell lines are MGMT (O-6-methylguanine-DNA methyltransferase) expressers and relatively resistant to the single chemotherapeutic agent [66,69] (Figure 5). Clinical trials are ongoing in advanced Ewing’s Sarcoma for safety and efficacy.

Vigil®

Vigil vaccine is a DNA engineered autologous whole tumor cell





Figure 6: Vigil™ is a plasmid of a bi-functional shRNA-furin DNA sequence which prevents cleavage of pro-TGFβ precursor into functional TGFβ1 & TGFβ2, as well as a GMCSF DNA sequence which stimulates MHC1 expression, antigen presentation and adaptive immune response. Autologous tumor cells are transfected with the plasmid via electroporation providing full tumor antigen profile. Prior studies demonstrated in Phase I, II testing induction of circulating cytotoxic T lymphocytes with increased immune response to tumor cells, measured by IFNγ ELISPOT response. Locally injected transfected autologous tumor cells increasingly express antigens to attracted dendritic cells, while immunosuppressive cytokines (such as TGFβ) are blocking tumor-induced immune tolerance and escape. Antigen-loaded dendritic cells activate naive CD8+ T cells in primary lymphatic tissue and enhance activation of cytotoxic CD8+ effector T cells that are then enabled to circulate to target lesions for cancer-antigen specific immune response. (Ghisoli, Barve et al. 2016).

immunotherapy which activates the afferent arm of the immune response arc by i) using autologous tumor cells as a source of the full matrix of tumor antigens, ii) recruiting, enhancing function and stimulating maturation of antigen-presenting cell (APC) populations via DNA encoded GMCSF expression and iii) dampening the escape of immune tolerance via knockdown of immunosuppressive TGFβ isoforms.

Vigil contains a plasmid comprised of both a DNA segment encoding for GMCSF protein expression and a bi-functional shRNA^{furin} DNA segment encoding for knockdown of Furin protein expression (a proprotein convertase which activates TGFβ1 and 2 isoforms) and consequent knockdown of both TGFβ1 and 2 (Figure 6).

In established cancers, TGFβ is an immune-suppressive cytokine, released by T-regulatory cells and cancer cells. Interestingly, it has paradoxical and context-dependent effects functioning as a tumor suppressor early in tumorigenesis and as an immune suppressive protein in the immune escape process and in established malignancies. In the latter context, TGFβ promotes cancer progression and proliferation, enhances activation of T-regulatory cells that contribute to apoptosis in APCs, and significantly decreases IFN-γ, granzymes A and B, and perforin release by cytotoxic T-cells [70]. By blocking these pathways, TGFβ suppresses the immune response and promotes immune-tolerance in cancer cells [71-74]. The immunogenic activity of plasmid-encoded, cell-secreted GMCSF has been extensively studied in a variety of GVAX trials and [gene-modified] oncolytic viral products [38,75-79].



The effective functionality of the Vigil encoded vectors is confirmed by product release criteria which require ≥ 30 pg of GMCSF secreted protein/ 10^6 cells and TGF β 1,2 knockdown of $\geq 30\%$ [78,80]. The gene modified autologous cells are irradiated to prevent tumor replication, placed in pharmaceutical standard vials containing 1×10^7 cells and then cryopreserved. The treatment protocol specifies intradermal administration of 1 mL vaccine every 4 weeks for ≥ 4 -12 doses depending on the quantity of vaccine available, Immune effectiveness is assessed by serial INF γ -ELISPOT analysis of PMBC response to autologous non-processed tumor cells as antigen source.

Phase I testing of Vigil in patients with advanced solid tumor types (heavily pre-treated and burdened with tumor volume) with evidence of progression following SOC therapies and/or FDA approved phase I/II clinical trials demonstrates safety and suggests efficacy of Vigil (1×10^7 cells or [in earlier trials] 2.5×10^7 cells per injection) with a correlation of survival to ELISPOT response (>10 spots/ 10^6 cells and 2x baseline) [78,80]. A long-term update of survival status of all Phase I treated patients [81] revealed a cohort of advanced Ewing's Sarcoma patients, predominantly third-line or greater, with suggestive evidence of survival benefit; i.e., $>75\%$ survival at 1 year compared to less than 25% survival based on historical experience. Longer term follow up of these patients also confirmed product safety with no evidence of Vigil related Grade ≥ 3 toxic effect.

More recently [82], a long term follow up of an expanded subset of advanced, late stage metastatic Ewing's Sarcoma patients treated with Vigil (n=16) was performed. The results of treatment in these patients were compared to the outcome of a non-randomized, concurrently treated group of Ewing's Sarcoma patients who underwent similar surgical procedures to harvest tissue for vaccine construction but who did not receive Vigil (n=14) for a variety of personal and/or physician determined reasons [82]. The median OS of Vigil treated patients (n=16) was 24 months vs. 6.8 months (Kaplan-Meier

in the control-group (no-Vigil treatment); a 17.2 month survival improvement (Figure 7) [82]. The update also showed a 75% 1-year survival of patients that received Vigil vs. 23% no-Vigil. Based on these findings, a randomized Phase IIb clinical trial in patients with advanced relapsed or refractory Ewing's Sarcoma who have not received \geq third-line therapy is active and ongoing.

Conclusion

Given historical lack of demonstrable effectiveness as well as a narrow therapeutic window, there are no FDA indicated treatment options for second- and third-line therapy of Ewing's Sarcoma patients who frequently have cumulative chemotherapy related toxic thereby limiting experimental treatment eligibility opportunity. As a result of advances in "-omics" analysis and molecular immunology as well as their directed application to Ewing's Sarcoma patients, a long needed window of opportunity has opened for exploration of innovative therapeutic options. Some evidence of activity has been suggested with single agent IGF-R1 and PARP inhibitors but, more importantly, even with failures data has been accrued and next generation studies have been implemented. Elements including biomarker identification, "-omic" analysis, pharmacokinetics and pharmacodynamics are now in place to help identify agent-specific sensitive subsets of Ewing's Sarcoma patients and guide protocol construction. Preclinical results with YK-4-279 and bi-shRNA EWS/FLI1 Type 1 LPX are encouraging, but on the basis of clinical results to date remain preliminary. Results from the phase I/II Vigil studies have matured and, based on analysis of outcomes, Vigil immunotherapy is currently undergoing randomized testing to determine qualification for FDA registration opportunity.

Acknowledgement

We gratefully acknowledge the generous support of the Alan B. Slifka Foundation, the Carson Sarcoma Foundation, the Chemo Warrior Foundation, Don and Linda Carter, The Crystal Charity Ball, the Jasper L. and Jack Denton Wilson Foundation, the Helen L. Kay Charitable Trust, Hyundai Hopes On Wheels, The Marilyn Augur Family Foundation, Michele Ashby, the Rutledge Foundation, the family and friends of Sam Day, the Speedway Children's Charities, Summerfield G. Roberts Foundation, Triumph Over Kid Cancer, Wipe Out Kids' Cancer and Young Texans Against Cancer. The authors would like to acknowledge Michelle Watkins and Brenda Marr for their competent and knowledgeable assistance in the preparation of the manuscript.

References

- Cotterill SJ, Ahrens S, Paulussen M, Jürgens HF, Voûte PA, Gadner H, et al. Prognostic factors in Ewing's tumor of bone: analysis of 975 patients from the European Intergroup Cooperative Ewing's Sarcoma Study Group. *J Clin Oncol.* 2000; 18: 3108-3114.
- Stahl M, Ranft A, Paulussen M, Bölling T, Vieth V, Bielack S, et al. Risk of recurrence and survival after relapse in patients with Ewing sarcoma. *Pediatr Blood Cancer.* 2011; 57: 549-553.
- Esiashvili N, Goodman M, Marcus RB Jr. Changes in incidence and survival of Ewing sarcoma patients over the past 3 decades: Surveillance Epidemiology and End Results data. *J Pediatr Hematol Oncol.* 2008; 30: 425-430.
- Downing J R, Head DR, Parham DM, Douglass EC, Hulshof MG, Link MP, et al. Detection of the (11;22) (q24;q12) translocation of Ewing's sarcoma and peripheral neuroectodermal tumor by reverse transcription polymerase chain reaction. *Am J Pathol.* 1993; 143: 1294-1300.

5. Aryee DN, Sommergruber W, Muehlbacher K, Dockhorn-Dworniczak B, Zoubek A, Kovar H, et al. Variability in gene expression patterns of Ewing tumor cell lines differing in EWS-FLI1 fusion type. *Lab Invest.* 2000; 80: 1833-1844.
6. Zoubek A, Dockhorn-Dworniczak B, Delattre O, Christiansen H, Niggli F, Gatterer-Menz I, et al. Does expression of different EWS chimeric transcripts define clinically distinct risk groups of Ewing tumor patients? *J Clin Oncol.* 1996; 14: 1245-1251.
7. de Alava E1, Kawai A, Healey JH, Fligman I, Meyers PA, Huvos AG, et al. EWS-FLI1 fusion transcript structure is an independent determinant of prognosis in Ewing's sarcoma. *J Clin Oncol.* 1998; 16: 1248-1255.
8. Le Deley MC, Delattre O, Schaefer KL, Burchill SA, Koehler G, Hogendoorn PC, et al. Impact of EWS-ETS fusion type on disease progression in Ewing's sarcoma/peripheral primitive neuroectodermal tumor: prospective results from the cooperative Euro-E.W.I.N.G. 99 trial. *J Clin Oncol.* 2010; 28: 1982-1988.
9. Parija T, Shirley S, Uma S, Rajalekshmy KR, Ayyappan S, Rajkumar T. Type 1 (11;22) (q24;q12) translocation is common in Ewing's sarcoma/peripheral neuroectodermal tumour in south Indian patients. *J Biosci.* 2005; 30: 371-376.
10. J Zucman, T Melot, C Desmaze, J Ghysdael, B Plougastel, M Peter, et al. Combinatorial generation of variable fusion proteins in the Ewing family of tumours. *EMBO J* 1993; 12: 4481-4487.
11. Sorensen PH, Lessnick SL, Lopez-Terrada D, Liu XF, Triche TJ, Denny CT. A second Ewing's sarcoma translocation, t(21;22), fuses the EWS gene to another ETS-family transcription factor, ERG. *Nat Genet.* 1994; 6: 146-151.
12. Peter M, Mugneret F, Aurias A, Thomas G, Magdelenat H, Delattre O. An EWS/ERG fusion with a truncated N-terminal domain of EWS in a Ewing's tumor. *Int J Cancer.* 1996; 67: 339-342.
13. Jeon IS, Davis JN, Braun BS, Sublett JE, Roussel MF, Denny CT, Shapiro DN. A variant Ewing's sarcoma translocation (7;22) fuses the EWS gene to the ETS gene ETV1. *Oncogene.* 1995; 10: 1229-1234.
14. Kaneko Y, Yoshida K, Handa M, Toyoda Y, Nishihira H, Tanaka Y, et al. Fusion of an ETS-family gene, EIAF, to EWS by t(17;22) (q12;q12) chromosome translocation in an undifferentiated sarcoma of infancy. *Genes Chromosomes Cancer.* 1996; 15: 115-121.
15. Urano F, Umezawa A, Hong W, Kikuchi H, Hata J. A novel chimera gene between EWS and E1A-F, encoding the adenovirus E1A enhancer-binding protein, in extraosseous Ewing's sarcoma. *Biochem Biophys Res Commun.* 1996; 219: 608-612.
16. Peter M, Couturier J, Pacquement H, Michon J, Thomas G, Magdelenat H, et al. A new member of the ETS family fused to EWS in Ewing tumors. *Oncogene.* 1997; 14: 1159-1164.
17. Antonescu CR. The role of genetic testing in soft tissue sarcoma. *Histopathology.* 2006; 48: 13-21.
18. Lazar A, Abruzzo LV, Pollock RE, Lee S, Czerniak B. Molecular diagnosis of sarcomas: chromosomal translocations in sarcomas. *Arch Pathol Lab Med.* 2006; 130: 1199-1207.
19. Leavey PJ, Collier AB. Ewing sarcoma: prognostic criteria, outcomes and future treatment. *Expert Rev Anticancer Ther.* 2008; 8: 617-624.
20. Bacci G, Ferrari S, Mercuri M, Longhi A, Giacomini S, Forni C, et al. Multimodal therapy for the treatment of nonmetastatic Ewing sarcoma of pelvis. *J Pediatr Hematol Oncol.* 2003; 25: 118-124.
21. Barker LM1, Pendergrass TW, Sanders JE, Hawkins DS. Survival after recurrence of Ewing's sarcoma family of tumors. *J Clin Oncol.* 2005; 23: 4354-4362.
22. Rasper M, Jabar S, Ranft A, Jürgens H, Amler S, Dirksen U, et al. The value of high-dose chemotherapy in patients with first relapsed Ewing sarcoma. *Pediatr Blood Cancer.* 2014; 61: 1382-1386.
23. Ferrari S, del Prever AB, Palmerini E, Staals E, Berta M, Balladelli A, et al. Response to high-dose ifosfamide in patients with advanced/recurrent Ewing sarcoma. *Pediatr Blood Cancer.* 2009; 52: 581-584.
24. Mora J, Cruz CO, Parareda A, de Torres C. Treatment of relapsed/refractory pediatric sarcomas with gemcitabine and docetaxel. *J Pediatr Hematol Oncol.* 2009; 31: 723-729.
25. Fox E, Patel S, Wathen JK, Schuetz S, Chawla S, Harmon D, et al. Phase II study of sequential gemcitabine followed by docetaxel for recurrent Ewing sarcoma, osteosarcoma, or unresectable or locally recurrent chondrosarcoma: results of Sarcoma Alliance for Research Through Collaboration Study 003. *Oncologist.* 2012; 17: 321.
26. Gaspar, ND, Hawkins S. Ewing Sarcoma: Current Management and Future Approaches Through Collaboration. *J Clin Oncol.* 2015; 33: 3036-3046.
27. Minas TZ, Han J, Javaheri T, Hong SH, Schleder M, Saygideğer-Kont Y, et al. YK-4-279 effectively antagonizes EWS-FLI1 induced leukemia in a transgenic mouse model. *Oncotarget.* 2015; 6: 37678-37694.
28. Erkizan HV, Schneider JA, Sajwan K, Graham GT, Griffin B, Chasovskikh S, et al. RNA helicase A activity is inhibited by oncogenic transcription factor EWS-FLI1. *Nucleic Acids Res.* 2015; 43: 1069-1080.
29. Erkizan HV, Kong Y, Merchant M, Schlottmann S, Barber-Rotenberg JS, Yuan L, et al. A small molecule blocking oncogenic protein EWS-FLI1 interaction with RNA helicase A inhibits growth of Ewing's sarcoma. *Nat Med.* 2009; 15: 750-756.
30. Barber-Rotenberg JS, Selvanathan SP, Kong Y, Erkizan HV, Snyder TM, Hong SP, et al. Single enantiomer of YK-4-279 demonstrates specificity in targeting the oncogene EWS-FLI1. *Oncotarget.* 2012; 3: 172-182.
31. Sung-Hyeok Hong, Sarah. E. Youbi, S. Peter Hong, Bhaskar Kallakury, Phillip Monroe, Hayriye V Erkizan, et al. Pharmacokinetic modeling optimizes inhibition of the 'undruggable' EWS-FLI1 transcription factor in Ewing Sarcoma. *Oncotarget.* 2014; 5: 338-350.
32. Toretsky JA, Erkizan V, Levenson A, Abaan OD, Parvin JD, Cripe TP, et al. Oncoprotein EWS-FLI1 activity is enhanced by RNA helicase A. *Cancer Res.* 2006; 66: 5574-5581.
33. Rahim S, Minas T, Hong SH, Justvig S, Çelik H, Kont YS, et al. A small molecule inhibitor of ETV1, YK-4-279, prevents prostate cancer growth and metastasis in a mouse xenograft model. *PLoS One.* 2014; 9: e114260.
34. Rao DD, Maples PB, Senzer N, Kumar P, Wang Z, Pappen BO, et al. Enhanced target gene knockdown by a bifunctional shRNA: a novel approach of RNA interference. *Cancer Gene Ther.* 2010; 17: 780-791.
35. Wang Z, Rao DD, Senzer N, Nemunaitis J. RNA interference and cancer therapy. *Pharm Res.* 2011; 28: 2983-2995.
36. Simmons O, Maples PB, Senzer N, Nemunaitis J. Ewing's Sarcoma: Development of RNA Interference-Based Therapy for Advanced Disease. *ISRN Oncol.* 2012; 247657.
37. Templeton NS, Lasic DD, Frederik PM, Strey HH, Roberts DD, Pavlakis GN. Improved DNA: liposome complexes for increased systemic delivery and gene expression. *Nat Biotechnol.* 1997; 15: 647-652.
38. Rao DD, Jay C, Wang Z, Luo X, Kumar P, Eysenbach H, et al. Preclinical Justification of pbi-shRNA EWS/FLI1 Lipoplex (LPX) Treatment for Ewing's Sarcoma. *Mol Ther.* 2016.
39. Prieur A, Tirode F, Cohen P, Delattre O. EWS/FLI-1 silencing and gene profiling of Ewing cells reveal downstream oncogenic pathways and a crucial role for repression of insulin-like growth factor binding protein 3. *Mol Cell Biol.* 2004; 24: 7275-7283.
40. Tirado OM, Mateo-Lozano S, Villar J, Dettin LE, Lloret A, Gallego S, et al. Caveolin-1 (CAV1) is a target of EWS/FLI-1 and a key determinant of the oncogenic phenotype and tumorigenicity of Ewing's sarcoma cells. *Cancer Res.* 2006; 66: 9937-9947.

41. Denduluri SK, Idowu O, Wang Z, Liao Z, Yan Z, Mohammed MK, et al. Insulin-like growth factor (IGF) signaling in tumorigenesis and the development of cancer drug resistance. *Genes Dis.* 2015; 2: 13-25.
42. Surmacz, E. Function of the IGF-I receptor in breast cancer. *J Mammary Gland Biol Neoplasia.* 2000; 5: 95-105.
43. Shawver LK, Slamon D, Ullrich A. Smart drugs: tyrosine kinase inhibitors in cancer therapy. *Cancer Cell.* 2002; 1: 117-123.
44. Surmacz, E. Growth factor receptors as therapeutic targets: strategies to inhibit the insulin-like growth factor I receptor. *Oncogene.* 2003; 22: 6589-6597.
45. Scotlandi K, Maini C, Manara MC, Benini S, Serra M, Cerisano V, et al. Effectiveness of insulin-like growth factor I receptor antisense strategy against Ewing's sarcoma cells. *Cancer Gene Ther.* 2002; 9: 296-307.
46. Strammiello R, Benini S, Manara MC, Perdichizzi S, Serra M, Spisni E, et al. Impact of IGF-I/IGF-IR circuit on the angiogenic properties of Ewing's sarcoma cells. *Horm Metab Res.* 2013; 35: 675-684.
47. Olmos D, Postel-Vinay S, Molife LR, Okuno SH, Schuetze SM, Paccagnella ML, et al. Safety, pharmacokinetics, and preliminary activity of the anti-IGF-1R antibody figitumumab (CP-751,871) in patients with sarcoma and Ewing's sarcoma: a phase I expansion cohort study. *Lancet Oncol.* 2010; 11: 129-135.
48. Juergens H, Daw NC, Geoerger B, Ferrari S, Villarroel M, Aerts I, et al. Preliminary efficacy of the anti-insulin-like growth factor type I receptor antibody figitumumab in patients with refractory Ewing sarcoma. *J Clin Oncol.* 2001; 29: 4534-4540.
49. Pappo AS, Patel SR, Crowley J, Reinke DK, Kuenkele KP, Chawla SP, et al. R1507, a monoclonal antibody to the insulin-like growth factor I receptor, in patients with recurrent or refractory Ewing sarcoma family of tumors: results of a phase II Sarcoma Alliance for Research through Collaboration study. *J Clin Oncol.* 2001; 29: 4541-4547.
50. Tap WD, Demetri G, Barnette P, Desai J, Kavan P, Tozer R, Benedetto PW, et al. Phase II study of ganitumab, a fully human anti-type-1 insulin-like growth factor receptor antibody, in patients with metastatic Ewing family tumors or desmoplastic small round cell tumors. *J Clin Oncol.* 2012; 30: 1849-1856.
51. Bhatia S, Krailo MD, Chen Z, Burden L, Askin FB, Dickman PS, et al. Therapy-related myelodysplasia and acute myeloid leukemia after Ewing sarcoma and primitive neuroectodermal tumor of bone: A report from the Children's Oncology Group. *Blood.* 2007; 109: 46-51.
52. Friedman DL, Whitton J, Leisenring W, Mertens AC, Hammond S, Stovall M, et al. Subsequent neoplasms in 5-year survivors of childhood cancer: the Childhood Cancer Survivor Study. *J Natl Cancer Inst.* 2010; 102: 1083-1095.
53. Rihani R, Bazzeh F, Faqih N, Sultan I. Secondary hematopoietic malignancies in survivors of childhood cancer: an analysis of 111 cases from the Surveillance, Epidemiology, and End Result-9 registry. *Cancer.* 2010; 116: 4385-4394.
54. Mulvihill MJ, Cooke A, Rosenfeld-Franklin M, Buck E, Foreman K, Landfair D, et al. Discovery of OSI-906: a selective and orally efficacious dual inhibitor of the IGF-1 receptor and insulin receptor. *Future Med Chem.* 2009; 1: 1153-1171.
55. Kim MY, Zhang T, Kraus WL. Poly (ADP-ribosyl)ation by PARP-1: 'PAR-laying' NAD⁺ into a nuclear signal. *Genes Dev.* 2005; 19: 1951-1967.
56. Gibson BA, Kraus WL. New insights into the molecular and cellular functions of poly (ADP-ribose) and PARPs. *Nat Rev Mol Cell Biol.* 2012; 13: 411-424.
57. Gibson BA, Kraus WL. Oral poly (ADP-ribose) polymerase inhibitor olaparib in patients with BRCA1 or BRCA2 mutations and recurrent ovarian cancer: a proof-of-concept trial. *Lancet.* 2010; 376: 245-251.
58. Bryant HE, Schultz N, Thomas HD, Parker KM, Flower D, Lopez E, et al. Specific killing of BRCA2-deficient tumours with inhibitors of poly(ADP-ribose) polymerase. *Nature.* 2005; 434: 913-917.
59. Balmaña J, Domchek SM, Tutt A, Garber JE. Stumbling blocks on the path to personalized medicine in breast cancer: the case of PARP inhibitors for BRCA1/2-associated cancers. *Cancer Discov.* 2001; 1: 29-34.
60. Lord, CJ, Ashworth A. The DNA damage response and cancer therapy. *Nature.* 2012; 481: 287-294.
61. Garnett MJ, Edelman EJ, Heidorn SJ, Greenman CD, Dastur A, Lau KW, et al. Systematic identification of genomic markers of drug sensitivity in cancer cells. *Nature.* 2012; 483: 570-575.
62. Brenner JC, Feng FY, Han S, Patel S, Goyal SV, Bou-Maroun LM, et al. PARP-1 inhibition as a targeted strategy to treat Ewing's sarcoma. *Cancer Res.* 2012; 72: 1608-1613.
63. Brohl AS, Solomon DA, Chang W, Wang J, Song Y, Sindiri S, et al. The genomic landscape of the Ewing Sarcoma family of tumors reveals recurrent STAG2 mutation. *PLoS Genet.* 2014; 10: e1004475.
64. Choy E, Butrynski JE, Harmon DC, Morgan JA, George S, Wagner AJ, et al. Phase II study of olaparib in patients with refractory Ewing sarcoma following failure of standard chemotherapy. *BMC Cancer.* 2014; 14: 813.
65. Murai J, Huang SY, Das BB, Renaud A, Zhang Y, Doroshow JH, et al. Trapping of PARP1 and PARP2 by Clinical PARP Inhibitors. *Cancer Res.* 2012; 72: 5588-5599.
66. Gill SJ, Travers J, Pshenichnaya I, Kogera FA, Barthorpe S, Mironenko T, et al. Combinations of PARP Inhibitors with Temozolomide Drive PARP1 Trapping and Apoptosis in Ewing's Sarcoma. *PLoS One.* 2015; 10: e0140988.
67. Engert F, Schneider C, Weiß LM, Probst M, Fulda S. PARP Inhibitors Sensitize Ewing Sarcoma Cells to Temozolomide-Induced Apoptosis via the Mitochondrial Pathway. *Mol Cancer Ther.* 2015; 14: 2818-2830.
68. Wilcoxon KM. The PARP inhibitor niraparib demonstrates synergy with chemotherapy in treatment of patient derived Ewing's sarcoma tumor Graft models. 2013.
69. Smith MA, Reynolds CP, Kang MH, Kolb EA, Gorlick R, Carol H, et al. Synergistic activity of PARP inhibition by talazoparib (BMN 673) with temozolomide in pediatric cancer models in the pediatric preclinical testing program. *Clin Cancer Res.* 2015; 21: 819-832.
70. Thomas, DA, Massague J. TGF-beta directly targets cytotoxic T cell functions during tumor evasion of immune surveillance. *Cancer Cell.* 2005; 8: 369-380.
71. Gorelik L, Flavell RA. Abrogation of TGFbeta signaling in T cells leads to spontaneous T cell differentiation and autoimmune disease. *Immunity.* 2000; 12: 171-181.
72. Gorelik LI, Flavell RA. Immune-mediated eradication of tumors through the blockade of transforming growth factor-beta signaling in T cells. *Nat Med.* 2001; 7: 1118-1122.
73. Massague, J. TGFbeta in Cancer. *Cell.* 2008; 134: 215-230.
74. Soares KC, Rucki AA, Kim V, Foley K, Solt S, Wolfgang CL, et al. TGF-beta blockade depletes T regulatory cells from metastatic pancreatic tumors in a vaccine dependent manner. *Oncotarget.* 2015; 6: 43005-43015.
75. Nemunaitis, J. Vaccines in cancer: GVAX, a GM-CSF gene vaccine. *Expert Rev Vaccines.* 2005; 4: 259-274.
76. Senzer NN, Kaufman HL, Amatruda T, Nemunaitis M, Reid T, Daniels G, Gonzalez R, et al. Phase II clinical trial of a granulocyte-macrophage colony-stimulating factor-encoding, second-generation oncolytic herpesvirus in patients with unresectable metastatic melanoma. *J Clin Oncol.* 2009; 27: 5763-5771.
77. Senzer, N., C. Bedell. OncoVEXGM-CSF: An Oncolytic Viral Immunotherapeutic in Melanoma. *Drugs Future.* 2010; 25: 449.

78. Senzer, N., M. Barve, et al. Long Term Follow Up: Phase I Trial of bi-shRNA furin/GMCSF DNA/Autologous Tumor Cell Immunotherapy (FANG™) in Advanced Cancer *Journal of Vaccines and Vaccination*. 2013; 4: 209.
79. Kaufman HL, Amatruda T, Reid T, Gonzalez R, Glaspy J, Whitman E, et al. Systemic versus local responses in melanoma patients treated with talimogene laherparepvec from a multi-institutional phase II study. *J Immunother Cancer*. 2016; 4: 12.
80. Senzer N, Barve M, Kuhn J, Melnyk A, Beitsch P, Lazar M, et al. Phase I trial of bi-shRNAi(furin)/GMCSF DNA/autologous tumor cell vaccine (FANG) in advanced cancer. *Mol Ther*. 2012; 20: 679-686.
81. Ghisoli M, Barve M, Schneider R, Mennel R, Lenarsky C, Wallraven G, et al. Pilot Trial of FANG Immunotherapy in Ewing's Sarcoma. *Mol Ther*. 2015; 23: 1103-1109.
82. Ghisoli M, Barve M, Mennel R, Lenarsky C, Horvath S, Wallraven G, et al. Three Year Follow up of GMCSF/bi-shRNAfurin DNA Transfected Autologous Tumor Immunotherapy (Vigil) in Metastatic Advanced Ewing's Sarcoma. *Mol Ther*. 2016.