



Chloroma- Myelosarcoma or Leukemic Nerve Tumor?

Grisold W*, Meng S and Grisold A

Department Neurology, Medical University of Vienna, Austria

Abstract

Following our review on leukemia and the nervous system (Grisold W) we would like to draw your attention to a rare manifestation of leukemia termed historically chloroma (CH) or myelosarcoma, which can rarely affect the cranial nerves and the peripheral nerves. These rare associations of leukemia and the peripheral nervous system, can occur in all stages of leukemia, as the first manifestation, as recurrence and post bone marrow transplant. Most frequently CH are seen in AML, less frequent CML, but also in OMF (?). They can also appear isolated, and leave room for speculations why leukemia can covert from a liquid into a solid tumor, and why the presentation can be in peripheral nerves. Chloromas, myelosarcomas or leukemic tumors are rare. Present at many sites of the body. Compared to lymphomas focal nerve lesion in leukemia are rare. For this review the presentation of CH in the the peripheral nervous system, including the cranial nerves was chosen. Peripheral nerve and CN tumors are rare. In leukemia the most frequent PNS lesions are meningeal involvement (LC), whereas neoplastic infiltration or compression of nerves by solid masses of leukemia are rare. Affection of CN and peripheral nerves is less frequently observed in leukemia, than in lymphoma.

Leptomeningeal Involvement or Not?

Meningeal spread of leukemia is not infrequent. In particular the triad between CNS, CN and radicular symptoms is characteristic. CH do not necessarily involve the meninges, however proximity of CH to the meninges either intracranially or spinal does not exclude an additional LC. The following considerations focus on CH.

Head and Cranial Nerves

CH can present in the orbit [1-12], infiltrate the optic nerve [13], affect the cavernous sinus [14] and causing focal CN damage, and also cause isolated hearing loss [15]. In addition mastoid or temporal bone lesions can cause CN dysfunction. CN can also be involved in leukemia in meningeal seeding (LC), dural involvement and also solid leukemic tumors presenting orbit, the optic nerve, in the cavernous sinus, with hearing loss, have been reported.

OPEN ACCESS

*Correspondence:

Grisold W, Department Neurology,
Medical University of Vienna, Austria,
E-mail: grisoldw@gmail.com

Received Date: 19 May 2017

Accepted Date: 13 Aug 2017

Published Date: 18 Aug 2017

Citation:

Grisold W, Meng S, Grisold A.
*Chloroma- Myelosarcoma or Leukemic
Nerve Tumor?. Clin Oncol. 2017; 2:
1329.*

Copyright © 2017 Grisold W. This is
an open access article distributed under
the Creative Commons Attribution
License, which permits unrestricted
use, distribution, and reproduction in
any medium, provided the original work
is properly cited.

Roots

The nerve roots can be a site of leukemic infiltration also presenting as a mass lesion. The involvement of the cauda equina by a leukemic mass has been observed [16-18].

Plexus

The brachial plexus as well as the lumbar and sacral plexus can be involved by CH. Due to modern imaging techniques the detection rate has improved, as the brachial, lumbar and sacral plexus can be easily examined. From the literature, the brachial plexus [19-23]. Seems more involved than the lumbar and sacral plexus. However this may also be an artifact due more specific symptom based evaluation of the brachial plexus.

The lumbar plexus can present with low back pain [24]. Also the sacrum can be the site CH with local nerve lesions [25-27].

Mononeuropathies

The presentation of CH in peripheral nerves presenting as mononeuropathies is rare. In some conditions also the term neuroleukemiosis has been suggested [28,29], Leukemia can infiltrate nerves or present as a diffuse parenchymatous tumor also affecting nerves [30], or presenting as a solid nerve tumor. On the UE both the median and the ulnar nerve have been described [31] and ulnar nerve [32]. In the lower extremities most cases have been observed in the femoral [33] and sciatic [34,35]. In the clinical setting several differential diagnoses ranging from entrapment

neuropathies, rare isolated effects of chemotherapy, and peripheral nerve tumors, as amyloidomas and rare leukemic deposition have to be considered. This is in particular difficult in cases in remission or after transplantation. In leukemia also coagulopathies can cause focal hemorrhages into peripheral nerves resulting in painful mononeuropathies.

Muscle

Also skeletal muscle can be the site of CHs [30], Muscle: massteric [6].

Discussion

The occurrence of a solid mass as presentation of leukemia (CH), is a rare event which can occur as a presentation, during the course of leukemia, as a relapse or as a complication of bone marrow transplant.

Imaging with MR, US and PET [17,11], has facilitated the detection, which may point of a more frequent appearance (However precise clinical criteria for the appearance of CH in ultrasound and MR are lacking.

The appearance of leukemia and CH in peripheral nerves is interesting. It has been proposed that the peripheral nerves and the peripheral nerve blood barrier may be an ideal situation to leukemic cells to survive in a sanctuary. However in most cases of leukemia the CN and peripheral nerves are spared from tumor infiltration, except the in intra - meningeal part, where LC is a frequent event. Another interesting aspect is the transformation of a liquid cancer into a solid cancer, at times isolated as a solid cancer, without the liquid presentation.

The appearance of CH is often uncharacteristic, and may require biopsy. Peripheral tumors are rare, the differential diagnostic approach needs to include several other differential diagnosis in particular if the CH is the first manifestation of leukemia. Once leukemia has been diagnosed this is more likely in acute and myeloid leukemia, and it has been described as recurrence or after BMT.

References

- Badreeldine A. Leucémie aiguë lymphoblastique révélée par une atteinte du sinus caverneux chez un jeune homme. *Pan Afr Med J.* 2014; 19: 98.
- Ala Alami B, Maaroufi M. Acute lymphoblastic leukemia revealed by an invasion of the cavernous sinus in a young man. *Pan Afr Med J.* 2014.
- Aregawi DG, Sherman JH, Douvas MG, Burns TM, Schiff D. Neuroleukemiosis: case report of leukemic nerve infiltration in acute lymphoblastic leukemia. *Muscle Nerve.* 2008; 38(3): 1196-1200.
- Bakst R, Jakubowski A, Yahalom J. Recurrent neurotropic chloroma: report of a case and review of the literature. *Adv Hematol.* 2011; 2011: 854240.
- Bakst R, Wolden S, Yahalom J. Radiation therapy for chloroma (granulocytic sarcoma). *Int J Radiat Oncol Biol Phys.* 2012; 82(5): 1816-1822.
- Bassichis B, McClay J, Wiatrak B. Chloroma of the masseteric muscle. *Int J Pediatr Otorhinolaryngol.* 2000; 53(1): 57-61.
- Boletto G, Miche M, Salam N, Eschard JP, Salmon JH. Low back pain and femoral neuralgia revealing myeloid sarcoma with megakaryocytic differentiation. *Joint Bone Spine.* 2016; 83(6): 743-744.
- Buakhao J, Tansawet A. Cauda equina involvement in acute myeloid leukemia relapse. *J Med Assoc Thai.* 2011; 94(10): 1271-1275.
- Zheng C, Liu X, Zhu W, Cai X, Wu J, Sun Z. Tailored central nervous system- directed treatment strategy for isolated CNS recurrence of adult acute myeloid leukemia. *Hematology.* 2014; 19(4): 208-212.
- Cavdar AO, Arcasoy A, Babacan E, Gözdaşoğlu S, Topuz U, Fraumeni JF Jr. Ocular granulocytic sarcoma (chloroma) with acute myelomonocytic leukemia in Turkish children. *Cancer.* 1978; 41(4): 1606-1609.
- Chhabra A, Thakkar RS, Andreisek G, Chalian M, Belzberg AJ, Blakeley J. Anatomic MR imaging and functional diffusion tensor imaging of peripheral nerve tumors and tumorlike conditions. *AJNR Am J Neuroradiol.* 2013; 34(4): 802-807.
- Eusebi V, Bondi A, Cancellieri A, Canedi L, Frizzera G. Primary malignant lymphoma of sciatic nerve. Report of a case. *Am J Surg Pathol.* 1990; 14(9): 881-885.
- Gokcan MK, Batikhan H, Calguner M, Tataragasi AI. Unilateral hearing loss as a presenting manifestation of granulocytic sarcoma (chloroma). *Otol Neurotol.* 2006; 27(1): 106-109.
- Grisold W, Briani C, Vass A. Malignant cell infiltration in the peripheral nervous system. *Handb Clin Neurol.* 2013; 115: 685-712.
- Grisold W, Briani C, Vass A. Leukemia and the Peripheral Nervous System. *Handb Clin Neurol.* 2013; 115: 685-712.
- Ha Y, Sung DH, Park Y, Kim du H. Brachial Plexopathy due to Myeloid Sarcoma in a Patient With Acute Myeloid Leukemia After Allogenic Peripheral Blood Stem Cell Transplantation. *Ann Rehabil Med.* 2013; 37(2): 280-285.
- Heckl S, Horger M, Faul C, Ebrahimi A, Ioanoviciu SD. Myeloid sarcoma of nervous plexus - infiltration of the nerve plexus by extramedullary manifestation of acute myeloid leukemia. *Rof.* 2014; 186(12): 1059-1062.
- Hurwitz BS, Sutherland JC, Walker MD. Central nervous system chloromas preceding acute leukemia by one year. *Neurology.* 1970; 20(8): 771-775.
- Onal IK, Shorbagi A, Göker H, Büyüksayık Y, Özçakar L, Tufan A, et al. Cauda equina syndrome as a rare manifestation of leukemia relapse during postallograft period. *J Natl Med Assoc.* 2006; 98(5): 808-810.
- Imrie KR, Kovacs MJ, Selby D, Lipton J, Patterson BJ, Pantalony D, et al. Isolated chloroma: the effect of early antileukemic therapy. *Ann Intern Med.* 1995; 123(5): 351-353.
- Kumar J, Seith A, Bakhshi S, Kumar R, Kumar A, Sen S. Isolated granulocytic sarcoma of the orbit. *Eur J Haematol.* 2007; 78(5): 456.
- Massoud M, Del Bufalo F, Caterina Musolino AM, Schingo PM, Gaspari S, Pisani M. Myeloid Sarcoma Presenting as Low Back Pain in the Pediatric Emergency Department. *J Emerg Med.* 2016.
- Mauermann ML, Angius D, Spinner RJ, Letendre LJ, Amrami KK, Dyck PJ. Isolated granulocytic sarcoma presenting as a brachial plexopathy. *J Peripher Nerv Syst.* 2008; 13(2): 153-156.
- Mosch A, Kuiters RR, Kazzaz BA. Intradural granulocytic sarcoma: a rare cause of sciatic pain. *Clin Neurol Neurosurg.* 1991; 93(4): 341-344.
- Novick SL, TL Nicol, EK Fishman. Granulocytic sarcoma (chloroma) of the sacrum: initial manifestation of leukemia. *Skeletal Radiol.* 1998; 27(2): 112-114.
- Paydas S, Zorludemir S, Ergin M. Granulocytic sarcoma: 32 cases and review of the literature. *Leuk Lymphoma.* 2006; 47(12): 2527-2541.
- Pochedly C. Neurologic manifestations in acute leukemia. III. Peripheral neuropathy and chloroma. *N Y State J Med.* 1975; 75(6): 878-882.
- Shah P, Yohendran J, Lowe D, McCluskey P. McCluskey. Devastating bilateral optic nerve leukaemic infiltration. *Clin Exp Ophthalmol.* 2012; 40(1): 114-115.
- Smith TR, Slimack N, McClendon J Jr, Wong A, Fessler RG. Low back pain and lumbar radiculopathy as harbingers of acute myeloid leukemia recurrence in a patient with myeloid sarcoma. *J Clin Neurosci.* 2012; 19(7): 1040-1041.

30. Song-Mee Cho, WH J. Granulocytic Sarcoma in the Leg Mimicking Hemorrhagic Abscess. JKSMRM. 2009; 13: 88-92.
31. Stillman MJ, Christensen W, Payne R, Foley KM. Leukemic relapse presenting as sciatic nerve involvement by chloroma (granulocytic sarcoma). Cancer. 1988; 62(9): 2047-2050.
32. Stork JT, Cigtay OS, Schellinger D, Jacobson RJ. Recurrent chloromas in acute myelogenous leukemia. AJR Am J Roentgenol. 1984; 142(4): 777-778.
33. Wang T, Miao Y, Meng Y, Li A. Isolated leukemic infiltration of peripheral nervous system. Muscle Nerve. 2015; 51(2): 290-293.
34. Warme B, Sullivan J, Tigrani DY, Fred DM. Chloroma of the forearm: a case report of leukemia recurrence presenting with compression neuropathy and tenosynovitis. Iowa Orthop J. 2009; 29: 114-116.
35. Yumi Ha M, Duk Hyun Sung, Yoonhong Park, Du Hwan Kim. Brachial Plexopathy due to Myeloid Sarcoma in a Patient With Acute Myeloid Leukemia After Allogenic Peripheral Blood Stem Cell Transplantation. Ann Rehabil Med. 2013; 37(2): 280-285.