



Perspective: Imaging for Gliomas

Yoshimasa Mori*

Department of Radiology and Radiation Oncology, Aichi Medical University, Japan

Perspective

The 2016 World Health Organization (WHO) classification of tumors of the central nervous system (CNS) broadly employs genetic alterations for diagnostic criteria, including isocitrate dehydrogenase-1 (IDH1) mutation or IDH2 mutation and 1p/19q codeletion [1], with the goal of creating more homogenous disease categories with greater prognostic value [2-5]. Molecular diagnostics is becoming an increasingly important aspect of clinical oncologic neuropathology practice. However, genetic analyses are time-consuming and costly. It is hoped that further accumulation of research work will help to clarify the correlations and discrepancies between the findings of molecular testing, conventional microscopic morphological examination of hematoxylin and eosin (HE) staining, and diagnostic imaging in clinical practice, despite potential conflicts [2]. Some investigations have tried to clarify the correlation between preoperative imaging and genetic status. Leu et al [6]. Reported that the combination of perfusion and diffusion-weighted magnetic resonance imaging (MRI), especially the combination of regional cerebral blood volume, apparent diffusion coefficient (ADC), T2 hyperintense volume, and presence of contrast enhancement may aid in non-invasively identifying genetic subtypes of diffuse gliomas. Brendle et al [7]. evaluated perfusion, dynamic contrast enhanced (DCE) perfusion imaging and arterial spine labeling (ASL). DCE and ASL perfusion were reported to be complementary in the differentiation of gliomas.

Genetic analyses are time-consuming and costly. More time is necessary than before between the acquisition of specimens at the surgical intervention and completion of their investigation including genetic alterations. The decision-making regarding the indications for placement of a nitrosourea wafer on the surgical operative field to cover the interval between the surgical intervention and beginning of drug therapy based on the definite diagnosis of malignant glioma, usually has to be made based on the preoperative imaging findings and quick histological diagnosis of frozen sections. It is hoped that a new regimen can be devised that will be able to considerably shorten this interval. It is also hoped that the optimal integration of genetic features, surface markers, HE histological morphological characteristics, and imaging findings including promising molecular and functional imaging will allow appreciable cost savings [8].

Arai et al. [9]. presented a case of glioblastoma, with atypical findings on MRI, arising during treatment with lenvatinib for thyroid cancer. MRI revealed a slightly high intensity lesion in the left frontal base area on T2-weighted or fluid attenuated inversion recovery (FLAIR) images, and the lesion showed only faint enhancement on T1-weighted images, after gadolinium administration. Though the histopathological diagnosis was glioblastoma, grade IV histology was observed in only a limited area and the majority of the specimen showed lower grade histology. They warned that molecular targeting agents used for other preexisting neoplasms might modify the imaging findings of brain tumors. The number of patients treated with chronic anti-neoplastic drugs is increasing. Opportunities to encounter such patients who developed brain tumors while receiving molecular targeting drugs are also likely increasing. In such patients not only the imaging findings but also the clinical course may be affected. They described that in patients receiving molecular target therapy for other diseases, it is necessary to consider a more aggressive pathology than indicated by the images. It is well known that anti-angiogenic agents often produce a rapid decrease in contrast enhancement because of reduced vascular permeability despite not having a true antitumor effect. It is always difficult to distinguish a decrease in contrast enhancement due to reduced vascular permeability by anti-angiogenic agents from true antitumor effects. Actually bevacizumab is used not only for its antitumor effects but also for anti-angiogenic ones to reduce peritumoral brain parenchymal edema and improve patient symptoms in the post radiotherapy period [10]. Furthermore, pseudoprogression is sometimes clinically observed during follow-up after chemotherapy for gliomas and radiation necrosis after radiotherapy, both of which would show enhancement by contrast medium on imaging. Besides at the initial diagnosis of diffuse glioma, during evaluation of the treatment results, for example associated with RANO criteria [11] after chemotherapy, particular

OPEN ACCESS

*Correspondence:

Yoshimasa Mori, Department of Radiology and Radiation Oncology, Aichi Medical University, Japan, Tel: 81-561-62-3311; Fax: 81-561-63-3268; E-mail: yoshim@aichi-med-u.ac.jp

Received Date: 14 Sep 2017

Accepted Date: 15 Nov 2017

Published Date: 24 Nov 2017

Citation:

Mori Y. Perspective: Imaging for Gliomas. Clin Oncol. 2017; 2: 1371.

Copyright © 2017 Yoshimasa Mori. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

care has to be taken in observing the follow-up imaging findings.

Modification of the imaging findings by previously administered drugs may be problematic in radiotherapy for glial tumors. During treatment planning of recent intensity modulated radiation therapy (IMRT) for glial tumors, pseudo-normalization of images, resulting in underestimation of tumor volume, may interfere with the appropriate delineation of the target volumes. Target volumes would be defined based on the gadolinium enhancement area and surrounding T2-weighted or FLAIR high-intense area in IMRT with simultaneous integrated boost (SIB) [12]. For example, Iuchi et al [13] employ 3-layered planning target volumes (PTVs) in a hypofractionated schedule. PTV1 was the surgical cavity and residual tumor on T1-weighted magnetic resonance images with 5-mm margins, PTV2 was the area with 15-mm margins surrounding the PTV1, and PTV3 was the high-intensity area on FLAIR images. Irradiation was performed in 8 fractions at total doses of 68, 40, and 32 Gy for PTV1, PTV2, and PTV3, respectively. They used concurrent temozolomide (TMZ) given at 75 mg/sq m/day for 42 consecutive days. Adjuvant TMZ was given at 150 to 200 mg/sq m/day for 5 days every 28 days. We also perform IMRT with SIB using 3 PTVs with a more conventional fraction schedule by TrueBeamSTx (Varian, Tokyo) linear accelerator with an ExacTrac (BrainLAB, Tokyo) patient positioning system. PTV1 is the surgical cavity and residual tumor on T1-weighted magnetic resonance images with 2-mm margins, PTV2 is the high-intensity area on preoperative fluid-attenuated inversion recovery images with 2-mm margins, and PTV3 is the area surrounding the PTV2 with 15-mm margins. Irradiation was performed in 30 fractions at total doses of 48 to 54, 60, and 69 Gy for PTV1, PTV2, and PTV3, respectively (unpublished data). We also use concurrent and adjuvant TMZ. Planning of these IMRT depends entirely on the preoperative and postoperative MRI findings.

Understanding is achieved by higher-detailed research datum accumulation of the associations among genetic features, surface markers, HE histological morphological characteristics, and imaging findings including molecular imaging and contrast enhancement depending on blood-brain-barrier conditions affected by not only tumor infiltration but also drug-induced alterations. It is hoped that greater advances in both molecular imaging and functional imaging will aid in the identification of the genetic characteristics affecting prognosis.

References

1. Louis DN, Wiestler OD, Cavenee WK. WHO Classification of Tumours of the Central Nervous System. International Agency for Research on Cancer. 2016.
2. Ballester LY, Huse JT, Tang G. Molecular Classification of Adult Diffuse Gliomas: Conflicting IDH1/IDH2, ATRX and 1p/19q Results. *Hum Pathol*. 2017.
3. Cimino PJ, Zager M, McFerrin L, Wirsching HG, Bolouri H, Hentschel B, et al. Multidimensional scaling of diffuse gliomas: application to the 2016 World Health Organization classification system with prognostically relevant molecular subtype discovery. *Acta Neuropathol Commun*. 2017; 22; 5(1):39.
4. Fuller CE, Jones DTW, Kieran MW. New classification for central nervous system tumors: Implications for diagnosis and therapy. *Am Soc ClinOncol Educ Book*. 2017; 37: 753-763.
5. Komori T. The 2016 WHO Classification of tumours of the central nervous system: The major points of revision. *Neurol Med Chir (Tokyo)*. 2017; 57: 301-311.
6. Leu K, Ott GA, Lai A, Nghiemphu PL, Pope WB, Yong WH, et al. Perfusion and diffusion MRI signatures in histologic and genetic subtypes of WHO grade II-III diffuse gliomas. *J Neurooncol*. 2017; 134: 177-188.
7. Brendle C, Hempel JM, Schittenhelm J, Skardelly M, Tabatabai G, Bender B, et al. Glioma grading and determination of IDH mutation status and ATRX loss by DCE and ASL perfusion. *J Neuroradiol*. 2017.
8. Smits M, van den Bent MJ. Imaging correlates of adult glioma genotypes. *Radiology*. 2017; 284: 316-331.
9. Arai N, Sasaki H, Tamura R, Ohara K, Yoshida K. Unusual magnetic resonance imaging findings of a glioblastoma arising during treatment with lenvatinib for thyroid cancer. *World Neurosurg*. 2017; 107: 1047.
10. Furuse M, Nonoguchi N, Kawabata S. Delayed brain radiation necrosis: pathological review and new molecular targets for treatment. *Review. Med MolMorphol*. 2015; 48(4): 183-90.
11. Eisele SC, Wen PY, Lee EQ. Assessment of brain tumor response: RANO and its offspring. *Curr Treat Options Oncol*. 2016; 17(7): 35.
12. Solberg TD, Agazaryan N, Goss BW. A feasibility study of 18F-fluorodeoxyglucose positron emission tomography targeting and simultaneous integrated boost for intensity-modulated radiosurgery and radiotherapy. *J Neurosurg*. 2004; 101 (3): 381-389.
13. Iuchi T, Hatano K, Kodama T, Sakaida T, Yokoi S, Kawasaki K, et al. Phase 2 trial of hypofractionated high-dose intensity modulated radiation therapy with concurrent and adjuvant temozolomide for newly diagnosed glioblastoma. *Int J Radiat Oncol Biol Phys*. 2014; 88(4): 793-800.