



# Potential Antioxidant Adjuvant Therapies in Breastcancer Treatments

Kuban-Jankowska A\*

Department of Medical Chemistry, Medical University of Gdansk, Poland

## Keywords

Cancer; Breast cancer; Adjuvant therapy; Lipoic acid; Resveratrol; Omega-3 fatty acids

## Editorial

Adjuvant therapy is an approach to fight with cancer as a support for standard anti-tumor therapy. Standard treatment includes combination of surgery, chemotherapy and radiation. The increasing resistance of cancer cells to chemotherapy revealed growing need of utilization of the supportive therapy in cancer treatment. Resistance of cancer cells occurs when they develop the ability to keep the chemotherapy drug from entering it or reduces the amount of the drug compounds that can enter to a level that does not cause damage. It is possible due to cancer cells produce a substance called p-glycoprotein, which can remove the chemotherapy drug from the cell. Many anticancer drugs require metabolic activation and thus cancer cells can develop resistance through decreased drug activation [1]. It was found that breast cancer cells can generate drug resistance [2]. In order to counteract resistance, chemotherapy drugs are usually administered as a combination of anti-cancer compounds, in case the cancer cells will be resistant to one of the drugs in the combination. Unfortunately, once cancer cells have developed resistance to one anti-cancer compound, it is more likely to develop resistance also to other type of drugs, making treatment more difficult. Thus it is so important to select the best possible drug combination and to use it as soon as the probability of resistance is at the lowest level. The main goal of supportive therapy in clinical practice is to induce cancer cells to be more sensitive to standard chemotherapy. Lipoic Acid (LA) is one of the potential natural compounds that can be considered as a supportive therapy in breast cancer. Numerous recent studies confirmed that LA can inhibit cell proliferation or increase apoptosis of, among others, breast, thyroid, lung, leukemia and colon cancer cells and suggested the possibility of lipoic acid as a potential anticancer agent [3-9]. It has been shown that lipoic acid and its analogs may have a significant impact in cancer metabolism and can be considered in chemotherapy [10]. It has been also discovered that anti-proliferative action of lipoic acid can be possibly due to the activation of pyruvate dehydrogenase, thus inducing the suppression of aerobic glycolysis and induction of cell death [11]. LA is a short-chain fatty acid, an organo sulfur compound derived from octanoic acid and can be synthesized in the human body, but it can be also absorbed from the diet. Recent studies have confirmed that lipoic acid utilized as a dietary supplement is well tolerated and reveals no significant toxicity [12]. Antioxidant compounds such as lipoic acid can be considered as a supplement to standard anticancer treatment to enhance the inhibitory effect on tumor cells and to protect from Reactive Oxygen Species (ROS) generated by anti-cancer agents. It was found that utilization of multiple antitumor drugs is associated with the induction of ROS [13]. Alpha-Lipoic Acid (ALA), also known as thioctic acid, can be reduced to dihydro lipoic acid (DHLA). Both compounds, ALA and its reduced form DHLA, are powerful antioxidants [14]. Other antioxidants that can be used as a supplementation of standard chemotherapy are, among others, resveratrol and omega-3 fatty acids. Resveratrol is a bioflavonoid found in many plants, mostly in grapes. Red wine is the main source of resveratrol in human diet. It was already many years ago discovered and confirmed nowadays that resveratrol can inhibit the growth of estrogen receptor positive human breast cancer cells [15-17]. Omega-3 are essential polyunsaturated fatty acids necessary for human health that must be delivered from diet. Omega-3 fatty acids, such as Eicosapentaenoic (EPA) and Docosahexaenoic (DHA) Acids are found in deep water fishes, among others, in mackerel, tuna and salmon [18]. Many studies pointed to the potential value of omega-3 fatty acids as adjuvant therapy to standard chemotherapy, as it has for example enhanced the efficiency of doxorubicin and mitomycin C in breast cancer cells [19,20]. The protective role of omega-3 fatty acids enriched diet was evidenced also for prostate and colon cancer [21,22].

## OPEN ACCESS

### \*Correspondence:

A Kuban-Jankowska, Department of Medical Chemistry, Medical University of Gdansk, Debinki 1 Street, 80210 Gdansk, Poland,

E-mail: alicjakuban@gumed.edu.pl

Received Date: 23 Mar 2018

Accepted Date: 20 May 2018

Published Date: 01 Jun 2018

### Citation:

Kuban-Jankowska A. Potential Antioxidant Adjuvant Therapies in Breastcancer Treatments. *Clin Oncol.* 2018; 3: 1483.

Copyright © 2018 Kuban-Jankowska A. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

## Conclusion

Antioxidants used as a supportive therapy, apart from the inhibitory properties against tumor cells by increasing their sensitivity to chemotherapy, may be also considered as the protection of healthy cells against free radicals and ROS produced by systemic anti-tumor drugs.

## Acknowledgements

This work was supported by Iuventus Plus Grant No. IP 2015 038 774 from Polish Ministry of Science and Higher Education.

## References

- Housman G, Byler S, Heerboth S, Lapinska K, Longacre M, Snyder N, et al. Drug Resistance in Cancer: An Overview. *Cancers (Basel)*. 2014;6(3):1769-92.
- Sengodan SK, K H S, Nadhan R, Srinivas P. Regulation of epithelial to mesenchymal transition by BRCA1 in breast cancer. *Crit Rev Oncol Hematol*. 2018;123:74-82.
- Li BJ, Hao XY, Ren GH, Gong Y. Effect of lipoic acid combined with paclitaxel on breast cancer cells. *Genet Mol Res*. 2015;14:17934-40.
- Jeon MJ, Kim WG, Lim S, Choi HJ, Sim S, Kim TY, et al. Alpha lipoic acid inhibits proliferation and epithelial mesenchymal transition of thyroid cancer cells. *Mol Cel Endocrinol*. 2016;419:113-23.
- Damnajanovic I, Kocic G, Najman S, Stojanovic S. Chemo preventive potential of alpha lipoic acid in the treatment of colon and cervix cancer cell lines. *Bratisl Lek Listy*. 2014;115(10):611-16.
- Kono Y, Inomata M, Hagiwara S, Hiratsuka T, Suzuki K, Koga H, et al. Anti proliferative effects of a new  $\alpha$ -lipoic acid derivative, DHL-HisZnNa, in HT29 human colon cancer cells in vitro. *Expert Opin Ther Targets*. 2012;16:103-09.
- Puchsaka P, Chaotham C, Chanvorachote P.  $\alpha$ -Lipoic acid sensitizes lung cancer cells to chemotherapeutic agents and anoikis via integrin  $\beta$ 1/ $\beta$ 3 down regulation. *Int J Oncol*. 2016;49(4):1445-56.
- Alpay M, Yurdakok-Dikmen B, Kismali G, Sel T. Anti leukemic effects of piperlongumine and alpha lipoic acid combination on Jurkat, MEC1 and NB4 cells in vitro. *J Cancer Res Ther*. 2016;12(2):556-60.
- Kuban-Jankowska A, Gorska-Ponikowska M, Wozniak M. Lipoic Acid Decreases the Viability of Breast Cancer Cells and Activity of PTP1B and SHP2. *Anticancer Res*. 2017;37(6):2893-98.
- Bingham PM, Stuart SD, Zachar Z. Lipoic acid and lipoic acid analogs in cancer metabolism and chemotherapy. *Expert Rev Clin Pharmacol*. 2014;7:837-46.
- Feuerecker B, Pirsig S, Seidl C, Aichler M, Feuchtinger A, Bruchelt G, et al. Lipoic acid inhibits cell proliferation of tumor cells in vitro and in vivo. *Cancer Biol Ther*. 2012;139(14):1425-35.
- Desideri I, Francolini G, Becherini C, Terziani F, DelliPaoli C, Olmetto E, et al. Use of an alpha lipoic, methylsulfonylmethane and bromelain dietary supplement (Opera<sup>®</sup>) for chemotherapy-induced peripheral neuropathy management, a prospectivestudy. *Med Oncol*. 2017;34(3):46.
- Carlisi D, Buttitta G, Di Fiore R, Scerri C, Drago-Ferrante R, Vento R, et al. Parthenolide and DMAPT exert cytotoxic effects on breast cancer stem-like cells by inducing oxidative stress, mitochondrial dysfunction and necrosis. *Cell Death Dis*. 2016;14(7):e2194.
- Biewenga GP, Haenen GR, Bast A. The pharmacology of the antioxidant lipoic acid. *Gen Pharmacol*. 1997;29(3):315-31.
- Lu R, Serrero G. Resveratrol, a natural product derived from grape, exhibits antiestrogenic activity and inhibits the growth of human breast cancer cells. *J Cell Physiol*. 1999;179(3):297-304.
- Suh J, Kim DH, Surh YJ. Resveratrol suppresses migration, invasion and stemness of human breast cancer cells by interfering with tumor-stromal cross-talk. *Arch Biochem Biophys*. 2018;643:62-71.
- da Costa Araldi IC, Bordin FPR, Cadoná FC, Barbisan F, Azzolin VF, Teixeira CF, et al. The in vitro radio sensitizer potential of resveratrol on MCF-7 breast cancer cells. *Chem Biol Interact*. 2018;282:85-92.
- Sargent JR. Fish oils and human diet. *Br J Nutr*. 1997;78(1):5-13.
- Larsson SC, Kumlin M, Ingelman-Sundberg M, Wolk A. Dietary long-chain n-3 fatty acids for the prevention of cancer: a review of potential mechanisms. *Am J Clin Nutr*. 2004;79(6):935-45.
- Berquin IM, Edwards IJ, Chen YQ. Multi-targeted therapy of cancer by omega-3 fattyacids. *Cancer Lett*. 2008;269:363-77.
- Berquin IM, Min Y, Wu R, Wu J, Perry D, Cline JM, et al. Modulation of prostate cancer genetic risk by omega-3 and omega-6 fattyacids. *J Clin Invest*. 2007;117(7):1866-75.
- Chapkin RS, Seo J, McMurray DN, Lupton JR. Mechanisms by which docosahexaenoic acid and related fatty acids reduce colon cancer risk and inflammatory disorders of the intestine. *Chem Phys Lipids*. 2008;153(1):14-23.