



A Retrospective Clinico-Pathologic Analysis of 100 Breast Cancer Cases: Experience from a Tertiary Care Hospital of Coastal India

Abhishek Singh Nayyar*

Department of Oral Medicine and Radiology, Maharashtra University of Health Sciences, India

Abstract

Context and aim: Breast cancers are relatively rare among young women amounting up to 6% of all breast carcinoma cases reported in that age group. In younger age groups, breast cancers, though, have a more aggressive behavior and poorer outcome in comparison to patients in the elderly age groups. The aim of the present study was to evaluate the histo-pathological characteristics of breast carcinomas reported in patients lesser than 40 years of age.

Materials and methods: Surgical specimens of carcinoma breast received at the department of pathology in a tertiary care hospital for a period of 2 years from January 2014 to December 2015 were included in the study. The histo-pathological slides were reviewed and clinical data was collected from the archival records and compared.

Statistical analysis: Statistical analysis was done using IBM SPSS statistics 17 (Chicago, USA). Various tumor characteristics were correlated with the morphological features and the other clinico-pathologic data using Chi-square value (X^2), paired t-test and Fischer's test. $p < 0.05$ was considered to be statistically significant.

Results: Infiltrating Ductal Carcinoma-Not Otherwise Specified (IDC-NOS) was found to be the most common variant of breast carcinoma in the younger age groups followed by invasive lobular carcinoma and mucinous carcinoma while stage III was the commonest stage reported in the younger age groups as against the elderly age groups wherein stage II was found to be commoner ($p = 0.006$).

Conclusion: The results of the present study revealed a more aggressive and advanced stages of breast carcinomas cases reported in the younger age groups.

Keywords: Breast cancer; Tertiary care settings; Younger age groups; TNM staging; Histo-pathology

Introduction

The most common cancers diagnosed in women are the breast cancers. These account for the second most common cause of cancer-related deaths in the affected women. Breast cancer is a relatively rare occurrence in the young women amounting to up to 6% of all breast carcinoma cases reported in that age group [1]. In younger age groups, breast cancers, though, have a more aggressive behavior and poorer outcome in comparison to patients in the elderly age groups [2-5]. Recently, a phenomenal change has been noticed in the behavior of breast carcinoma cases wherein more of the young women have been found to be more commonly getting afflicted and being diagnosed with breast carcinomas [6-8]. The aim of the present study was to evaluate the histo-pathological characteristics of breast carcinomas reported in patients lesser than 40 years of age and compare those with those reported in the elderly age groups.

Materials and Methods

A retrospective study was conducted over a period of 2 years from January 2014 to December 2015 in department of pathology of a tertiary care hospital of South India wherein surgical specimens of carcinoma breast received at the department of pathology were included. The histo-pathological slides were reviewed and clinical data was collected from the archival records and compared. All data including the clinical presentation, pathological type and TNM staging of the tumor were compared between the two groups. Carcinoma-*in-situ*, male breast carcinomas and inadequate biopsy samples

OPEN ACCESS

*Correspondence:

Abhishek Singh Nayyar, Department of Oral Medicine and Radiology, Maharashtra University of Health Sciences, Nashik, Maharashtra, India, Tel: +91-98509 04067; E-mail: singhabhishekndls@gmail.com

Received Date: 16 Jan 2020

Accepted Date: 17 Feb 2020

Published Date: 25 Feb 2020

Citation:

Nayyar AS. A Retrospective Clinico-Pathologic Analysis of 100 Breast Cancer Cases: Experience from a Tertiary Care Hospital of Coastal India. Clin Oncol. 2020; 5: 1685.

Copyright © 2020 Abhishek Singh Nayyar. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Table 1: Tumor characteristics in breast carcinoma cases reported in the 2 age groups.

Tumor Characteristics	<40 years	>40 years	p-value
n (No. of cases)	20	80	
Histological Type			
Infiltrating Ductal carcinoma	17	69	p>0.05
Lobular carcinoma	2	6	
Papillary carcinoma	0	2	
Mucinous carcinoma	1	2	
Neuro-endocrine carcinoma	0	1	
Tumor size			
T1	2	4	p<0.017*
T2	9	51	
T3	9	23	
T4	0	2	
Nodal status			
N0	8	41	p<0.042*
N1	7	24	
N2	3	10	
N3	2	5	
TNM Staging			
Stage 1	2	4	p<0.006*
Stage 2	12	55	
Stage 3	6	20	
Stage 4	0	1	
Immuno-histochemistry			
ER/PR Positive, Her2 Negative	3 (15%)	14 (17.50%)	p<0.043*
ER/PR Positive, Her2 Positive	4 (20%)	21 (26.20%)	
ER/PR Negative, Her2 Negative	8 (40%)	15 (18.80%)	
ER/PR Negative, Her2 Positive	2 (10%)	14 (17.50%)	
Not done	3 (15%)	16 (20%)	

p<0.05* Statistically Significant

were excluded from the study. The study was subjected for ethical approval from the Institutional Ethical Committee and clearance was obtained before the start of the study.

Statistical Analysis

Statistical analysis was done using IBM SPSS statistics 17 (Chicago, USA). Various tumor characteristics were correlated with the morphological features and the other clinico-pathologic data using Chi-square value (χ^2), paired t-test and Fischer's test. p<0.05 was considered to be statistically significant.

Results

A total of hundred surgical specimens were evaluated out of which 20 cases were reported in patients below 40 years while 80 specimens were from cases above 40 years of age (Table 1). Infiltrating Ductal Carcinoma-Not Otherwise Specified (IDC-NOS) (Figures 1a,1b and 2a,2b) was found to be the most common variant of breast carcinoma in the younger age groups followed by invasive lobular carcinoma and mucinous carcinoma (Table 1/Graph 1). Also, tumor size of diameter >5 cm (T3) was found to be more common in the younger age groups while in the elderly age groups, tumor size of diameter 2




Figure 1a,1b: Tumor cells showing areas of fibrosis and calcification (H&E, 100x, 200x).





Figure 2a,2b: Tumor cells showing hemosiderin laden macrophages along with areas of fibrosis (H&E, 100x, 200x).



Graph 1: Various histological types of breast carcinomas in the younger and elderly age groups.

cm to 5 cm (T2) was found to be more common (p=0.017) (Table 1). Furthermore, younger patients had greater number of lymph nodes involved in the malignant process (Figure 3a-3c) which were found to be positive as compared to the elderly age groups (p=0.042) (Table 1). Immunohistochemistry revealed ER/PR positivity while Her2 negativity in 15% of the cases lesser than 40 years of age and 17.5% of the cases more than 40 years of age. On the contrary, ER/PR Positivity as well as Her2 positivity was reported in 20% of the cases lesser than 40 years of age and 26.2% of the cases more than 40 years of age. Likewise, ER/PR negativity with Her2 negativity was reported in 40% of the cases lesser than 40 years of age and 18.8% of the cases more than 40 years of age as against ER/PR negativity and Her2 positivity reported in 10% of the cases lesser than 40 years of age and 17.5% of the cases more than 40 years of age and the results were found to be statistically significant (p<0.043) (Table 1). Also, in the younger age groups, stage III was the commonest stage reported as against the elderly age groups wherein stage II was found to be commoner

Table 2: Clinico-pathologic data correlated with tumor characteristics in breast carcinoma cases reported in the 2 age groups in the previous studies.

Studies	n (No. of cases)		Most common histological type of the tumor		Predominant tumor grade and TNM staging	
	<40 years	>40 years	<40 years	>40 years	<40 years	>40 years
Manilal et al. [6]	100 (23.80%)	420 (72.20%)	IDC-NOS (87%)	IDC-NOS (81%)	T3 (43.10%) TNM Stage-III (54.80%)	T2 (41%) TNM Stage-II (47%)
Sidoni et al. [11]	50	50 (50 Randomly selected matched cases)	IDC-NOS (80%)	IDC-NOS (90%)	TNM Stage-III (58%)	TNM Stage-II (75%)
Zhang Q et al. [14]	181 (37%)	488 (63%)	IDC-NOS (77.30%)	IDC-NOS (84.40%)	T2 (69.30%) TNM Stage-III (49.20%)	T2 (77.70%) TNM Stage-II (52.20%)
Thapa et al. [16]	263 (27.90%)	681 (72.10%)	IDC-NOS (93.10%)	IDC-NOS (86%)	T3 (55.90%) TNM Stage-III (55.34%)	T2 (54.20%) TNM Stage-II (57.90%)
Thangjam et al. [18]	160 (31%)	347 (68%)	IDC-NOS (85.60%)	IDC-NOS (83.50%)	T3 (58.13%) TNM Stage-III (55.34%)	T2 (69.45%) TNM Stage-II (64%)
Present study	20 (20%)	80 (80%)	IDC-NOS (85%)	IDC-NOS (86.20%)	T3 (60%) TNM Stage-III (60%)	T2 (63%) TNM Stage-II (68.70%)




Figure 3a-3c: Lymph nodal tissue showing tumor cells with areas of fibrosis and hyalinization (H&E, 100x, 200x, 400x).



Graph 2: TNM staging of breast carcinomas in the younger and elderly age groups.

(p=0.006) (Table 2/Graph 2).

Discussion

Although breast carcinomas are recognized to occur more commonly in the post-menopausal age, they can occur in the relatively young age, too. In younger age groups, breast cancers, though, have a more aggressive behavior and poorer outcome in comparison to patients in the elderly age groups [9-11]. The incidence of breast cancers is less by 2% to 3% in the west as compared to the Indian population [12]. As per Mohanti BK et al. [7], in young Indian women, the incidence of breast cancers was found to be around 5.5% close to the incidence recorded in the study conducted by Das U et

al. [8] which reported an incidence of 8%. The results of the present study showed a relative incidence of 20% of the cases reported in the younger age groups as per the data retrieved from the archival records of the hospital set-up. Clinically and histo-pathologically, breast cancers in the young are different from that reported in the elderly age groups [13]. In the present study, too, the average age at presentation was found to be 64.5 years based on the archival records of the hospital which was similar to the previously reported studies [14-17]. The oldest patient reported was of 78 years of age in the present study as against the youngest patient being just 33 years of age. Lump in the breast was the most common presenting complaint (83%) followed by mastalgia (6%) similar to the common complaints reported in other studies. In addition, T3 was found to be more common in the younger age groups while in the elderly age groups, T2 was found to be more common (p=0.017). On histo-pathological examination, stage III was the commonest stage reported in the younger age groups as against the elderly age groups wherein stage II was found to be commoner (p=0.006). Similar results were found in various other studies conducted in the past including the one conducted by Manilal et al. [6], Sidoni et al [11], Thapa et al. [16] and Thangjam et al. [18] wherein stage III was the predominant stage reported in the younger women (<40 years) while stage II was the most common stage reported in the elderly age groups (>40 years). This was in relative contrast to the results of the study conducted by Zhang Q et al. [14] which reported stage II to be the predominant stage in both the younger and elderly age groups. As per Colleoni M et al. [19], more aggressive behavior of the disease was seen in 48% of the young patients. In the present study, too, young patients had advanced stage of the disease with concomitant higher rate of lymph node positivity reported and greater number of lymph nodes involved

in the malignant process (25%) in the younger age groups as compared to the elderly patients (6.2%) ($p=0.042$). Also, IDC-NOS were found to be the most common variant of breast carcinoma reported in the younger age groups followed by invasive lobular carcinoma and mucinous carcinoma. This was, also, in agreement with the results of most of the studies conducted previously wherein IDC-NOS has been the most common histological type of breast carcinoma in both the younger and the elderly age groups [9-11,13-18]. Furthermore, the number of cases with triple negative breast carcinomas was found to be more in the younger age groups in the present study compared to the elderly age groups (40% vs. 18.8%) ($p<0.043$), a finding which was in agreement with most of the previous studies which reported triple negative breast carcinoma cases and high grades of tumor with lymph nodal involvement and poorer prognosis to be commonly seen in association with patients in the younger age groups [20-23].

Conclusion

Breast cancer diagnosed at young age has an aggressive behavior and poorer outcome in comparison to patients in the elderly age groups. Young women often tend to present with advanced stage of the disease and high grades of tumor with lymph nodal involvement and metastasis conferring an inferior prognosis. This illustrates the need for more clinical trials to be conducted on younger patients with breast cancers with an aim to improve the overall outcome in this age group. Breast cancer awareness and screening programs and easy access to health care delivery systems may help to increase the awareness and detect breast cancer in early stage. Thus, women in this age group must be educated for regular self examination and early medical attention on discovery of any breast lump that is persistent for more than certain duration. Also, clinical breast examination at health care delivery systems during routine check-ups should be encouraged to facilitate an early intervention if at all needed.

References

- Thomson CS, Brewster DH, Dewar JA, Twelves CJ; Scottish Cancer Therapy Network. Improvements in survival for women with breast cancer in Scotland between 1987 and 1993: Impact of earlier diagnosis and changes in treatment. *Eur J Cancer*. 2004;40(5):743-53.
- Peto R, Boreham J, Clarke M, Davies C, Beral V. UK and USA breast cancer deaths down 25% in year 2000 at ages 20-69 years. *Lancet*. 2000;355(9217):1822.
- Gnerlich JL, Deshpande AD, Jeffe DB, Sweet A, White N, Margenthaler JA. Elevated breast cancer mortality in women younger than age 40 years compared with older women is attributed to poorer survival in early stage disease. *J Am Col Surg*. 2009;208(3):341-7.
- El Saghir NS, Seoud M, Khalil MK, Charafeddine M, Salem ZK, Geara FB, et al. Effects of young age at presentation on survival in breast cancer. *BioMed Central Cancer*. 2006;6:194-7.
- Anders CK, Hsu DS, Broadwater G, Acharya CR, Foekens JA, Zhang Y, et al. Young age at diagnosis correlates with worse prognosis and defines a subset of breast cancers with shared patterns of gene expression. *J Clin Oncol*. 2008;26(20):3324-30.
- Manilal B, Bargain D, Patnayak R, Jena A, Reddy KS, Sampath V, et al. Breast cancer in young: Experience at a tertiary care center in Tirupati, Andhra Pradesh. *J Clin Sci Res*. 2016;5:93-100.
- Mohanti BK, Raina V, Gogia A, Deo SVS, Shukla NK. Young patients with breast cancer (<35 years): Single-institution study of 194 patients from India. *J Mol Biomark Diagn*. 2013;6:261-4.
- Das U, Lakshmaiah KC, Lokanatha D, Babu G, Jacob L, Babu S. Breast Cancer in Women of Younger than 35 Years: A Single Center Study. *J Mol Biomark Diagn*. 2015;6:261-4.
- Nixon AJ, Neuberger D, Hayes DF, Gelman R, Connolly JL, Schnitt S, et al. Relationship of patient age to pathologic features of the tumor and prognosis for patients with stage I or, II breast cancer. *J Clin Oncol*. 1994;12(5):888-94.
- Meng J, Lang R, Fan Y, Fu L. Clinico-pathological and biological features of breast cancer in young females and their relationship with prognosis. *Zhonghua Zhong Liu Za Zhi*. 2007;29(4):284-8.
- Sidoni A, Cavaliere A, Bellezza G, Scheibel M, Bucciarelli E. Breast cancer in young women: Clinico-pathological features and biological specificity. *Breast*. 2003;12(4):247-50.
- US Department of Health and Human Services. SEER Cancer Statistics Review: 1975-2014. National Cancer Institute. Bethesda, MD, USA.
- Walker RA, Lees E, Webb MB, Dearing SJ. Breast carcinomas occurring in young women (<35 years) are different. *Br J Cancer*. 1996;74(11):1796-800.
- Zhang Q, Ma B, Kang M. A retrospective comparative study of clinico-pathological features between young and elderly women with breast cancer. *Int J Clin Exp Med*. 2015;8(4):5869-75.
- Bertheau P, Steinberg SM, Cowan K, Merino MJ. Breast cancer in young women: Clinico-pathologic correlation. *Semin Diagn Pathol*. 1999;16(3):248-56.
- Thapa B, Singh Y, Sayami P, Shrestha UK, Sapkota R, Sayami G. Breast cancer in young women from a low risk population in Nepal. *Asian Pac J Cancer Prev*. 2013;14(9):5095-9.
- Raina V, Bhutani M, Bedi R, Sharma A, Deo SV, Shukla NK, et al. Clinical features and prognostic factors of early breast cancer at a major cancer center in North India. *Indian J Cancer*. 2005;42(1):40-5.
- Thangjam S, Laishram RS, Debnath K. Breast carcinoma in young females below the age of 40 years: A histo-pathological perspective. *South Asian J Cancer*. 2014;3(2):97-100.
- Colleoni M, Rotmensz N, Robertson C, Orlando L, Viale G, Renne G, et al. Very young women (<35 years) with operable breast cancer: Features of disease at presentation. *Ann Oncol*. 2002;13(2):273-9.
- Assi HA, Khoury KE, Dbouk H, Khalil LE, Mouhieddine TH, El Saghir NS. Epidemiology and prognosis of breast cancer in young women. *J Thorac Dis*. 2013;5(Suppl 1):S2-8.
- Fredholm H, Eaker S, Frisell J, Holmberg L, Fredriksson I, Lindman H. Breast cancer in young women: Poor survival despite intensive treatment. *PLoS One*. 2009;4(11):7695.
- Xiong Q, Valero V, Kau V, Kau SW, Taylor S, Smith TL, et al. Female patients with breast carcinoma age 30 years and younger have a poor prognosis: The MD Anderson Cancer Center experience. *Cancer*. 2001;92(10):2523-8.
- Chung M, Chang HR, Bland KI, Wanebo HJ. Younger women with breast carcinoma have a poorer prognosis than older women. *Cancer*. 1996;77(1):97-103.