



Socio-Demographic and Clinical Characteristics of Cancer Patients of Rangpur, Bangladesh

Ratindra Nath Mondal^{1*}, Shah¹, Sarwer Jahan², Mokhlesur Rahman Sarker², Jahan Afroza Khanum², Mahfuj-Ul-Anwar², Rahenur Mondol², Narayan Chandra Sarkar², Sajeda Afrin², Sara Jabeen Khandaker³ and Moni Ranil³

¹Department of Medicine, Society of General Physicians and Dakarkhana (GP Center), Bangladesh

²Department of Medicine, Rangpur Medical College, Bangladesh

³Department of Medicine, Rajshahi Medical College Hospital, Bangladesh

Abstract

Background: Cancer is the 6th leading cause of death and accounts for 10% of all mortality in Bangladesh. Main workforce structures in the country, age from 30 to 65 years are affected mostly.

Materials and Methods: This was a cross sectional study, carried out in Department of Oncology of Rangpur Medical College Hospital. Cancer patients attending in OPD were included in this study.

Results: We have studied a total of 313 cancer patients, male were more than female (59.7% vs. 40.3%), mean age was 48.92 years. 82.7% pt were from rural area, 54.6% were illiterate and monthly income less than 5,000 BDT in 85.9% cases. 54% of the patients were smoker and 8.3% patient used SLT (smokeless tobacco). Carcinoma of breast was the highest among all cancer 17.89%, which is followed by carcinoma of ovary 12.14%. But as a group majority of the cases were GIT cancer (oral cavity, tongue, esophagus, stomach and colon) 31.94% and respiratory tract cancer (larynx, lung) 23.64%. In male lung cancer (39.57%) and in female breast cancer (44.45%) was the most common cancer. Among the known risk factor of cancer-increasing age (mean age 48.92 years), smoking (54%) and radiation exposure (68%) were present. In our study, metastasis occurred at the time of diagnosis in 48.6% of the cases. 91.7% of the patients were suggested chemotherapy and the rest were both chemotherapy and radiotherapy.

Conclusion: Cancer is more common in older age, illiterate, poor socio-economic group and patient from rural area. Carcinoma of breast was the highest among all cancer. But as a group majority of the cases were GIT cancer. Approximately half of the cases were metastasized at the time of diagnosis.

Keywords: Cancer; Socio-demographic; Breast; Ovarian

Introduction

Cancer cases doubled globally between 1975 and 2000, will double again by 2020 and triple by 2030. The rapid increase in the global cancer burden represents a real challenge for health systems worldwide [1]. Globally, cancers in all forms are causing about 12% of all deaths. In developed countries cancer is the second leading cause of death accounting for 21% of mortality by other causes and in developing countries it ranks third, accounting for 9.5% of all deaths [2]. According to Bangladesh bureau of statistics cancer is the sixth leading cause of death in Bangladesh (BBS, 2004). International Agency for Research on Cancer (IARC) has estimated cancer-related death rate in Bangladesh to be 7.5% in 2005 and it will be increase to 13% in 2030 [3]. Recent studies conducted in Bangladesh revealed that nearly two million cancer patients currently exist in Bangladesh and every year about two hundred thousand patients are affected by cancer with death estimation of about 150,000 per year. At present, Bangladesh is known as the ninth most populous country in the world regarding cancer sufferings [4,5,6].

A WHO study estimates that there are 49,000 oral, 71,000 pharynx and laryngeal and 196,000 lung cancer cases in Bangladesh among those aged ≥ 30 years. The same study observed that 3.6% of the admissions in medical college hospitals for the same age group are due to cancers of oral cavity, larynx and lungs [3]. Although cancer is a devastating disease but it is largely preventable. If we have the adequate data about cancer patients, then by applying appropriate measures, a great impact on reducing the global cancer burden can be achieved. And one of the instruments for data collection of cancer patients is their registration [7]. Cancer registration is of two types-population

OPEN ACCESS

*Correspondence:

Ratindra Nath Mondal, Department of Medicine, Society of General Physicians and Dakarkhana (GP Center), Rangpur, Bangladesh, E-mail: dr.ratinmondal@gmail.com

Received Date: 17 Aug 2020

Accepted Date: 10 Sep 2020

Published Date: 12 Sep 2020

Citation:

Mondal RN, Shah, Jahan S, Sarker MR, Khanum JA, Mahfuj-Ul-Anwar, et al. Socio-Demographic and Clinical Characteristics of Cancer Patients of Rangpur, Bangladesh. *Clin Oncol.* 2020; 5: 1735.

Copyright © 2020 Ratindra Nath Mondal. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Based Cancer Registry (PBCR) and Hospital Based Cancer Registry (HBCR). HBCR limits its aim to record the particulars of cancer cases seen in a given hospital or group of hospitals irrespective of boundaries of geographical areas. HBCR records all cases in a given hospital, usually without knowledge of the background population; the emphasis is on clinical care, regular routine follow up and hospital administration. It may also form the nucleus for PBCR. Therefore this study was undergone to determine the socio-demographic patterns of cancer patients.

Materials and Methods

This was a cross sectional study, carried out in Department of Oncology of Rangpur Medical College Hospital, Bangladesh from 2nd May, 2017 to 1st November, 2017. Assuming unknown prevalence of cancer in this region a sample size of 384 was calculated to give the true prevalence with a precision of 5% with 95% of confidence level. On an average about 30 patient visit to Department of Oncology. From the daily visited patients every 3rd patient was taken to generate sample of 384. The first patient was chosen randomly from cancer patients 1 to 10.

Data collection

A non-medical staff was trained to collect the data, who is not informed about the details of the study. An informed consent was obtained before enrollment. Data regarding attributes of social classes; education of the patients, occupation and per capita income were collected by face to face interview. Medical records of the patients were taken and examined from the register of medical records of department of Oncology of Rangpur Medical College Hospital. After collection of data it was coded and checked manually, incomplete data sheets were removed; finally we had 313 data to be analyzed.

Statistical analysis

Data analysis was done according to the objectives of the study by using SPSS-17.0 (Statistical Program for Social Science) software program. The result of the clinical study and statistical analysis is presented in the form of text, table, bar and chart.

Ethical consideration

The ethical committee of Rangpur Medical College has approved the study protocol and questionnaire on 12th February, 2017.

Result

In this study, we have studied a total of 313 cancer patients of both sexes. Male were more than female (59.7% vs. 40.3%). Mean age of the study population was 48.92 years. (Table 1 shows the socio-demographic characteristics of the study population).

Fifty four percent (54%) of the patients were smoker, among them 48.2% gave up smoking after diagnosis of cancer but 5.8% continued smoking and 8.3% patient used SLT (smokeless tobacco) in the form of jorda and gul.

Carcinoma of breast was the highest among all cancer 17.89%, which is followed by carcinoma of ovary 12.14%. But as a group majority of the cases were GIT cancer (oral cavity, tongue, esophagus, stomach, colon) 31.94% and respiratory tract cancer (larynx, lung) 23.64%. Among the GIT cancer, carcinoma oral cavity 36% was the maximum followed by carcinoma of the stomach 28%, carcinoma of the esophagus 18% and carcinoma of the colon 18%. Carcinoma of the larynx, supraglottic region and bronchial carcinoma was 72.98% and 27.02% respectively in respiratory tract cancer group (Table 2).

Table 1: Socio-demographic characteristics of the study population (n=313).

Variables	Frequency	Percentage (%)
Mean age	48.92 years (SD 14.29) (age range 22 to 75 years)	
Sex Male	187	59.70%
Female	126	40.30%
Residence Rural	259	82.70%
Urban	54	17.30%
Level of education		
Illiterate	171	54.60%
<5 class	72	23%
5- <10 class	52	16.60%
>10	9	2.90%
Graduate and above	9	2.90%
Occupation		
Service	9	2.90%
Business	27	8.60%
Agriculture	133	42.50%
Housewife	117	37.40%
Others	27	8.60%
Monthly income		
<5000 taka*	269	85.90%
>5000 taka	44	14.10%

*1 dollar = 82 taka

Table 2: Frequency of different malignancy (n=313).

Name of cancer	Percentage/frequency
Carcinoma of breast	17.89% (56)
Carcinoma of ovary	12.14% (38)
GIT cancer (oral cavity, tongue, esophagus, stomach, colon)	31.94% (100)
Respiratory tract cancer (larynx, lung)	23.64% (74)
Carcinoma cervix	2.88% (9)
Carcinoma gall bladder	2.88% (9)
Carcinoma of neck lymph node with unidentified primary site	8.62% (27)

Table 3: Risk factors of cancer.

Risk factors	Percentage (frequency)
Age	48.92 years (SD 14.29) (age range 22 to 75 years)
Smoking	54%
History of taking added salt	51.80%
Radiation exposure	68.70%
Sun exposure	80.20%
Obesity	1.90%
Over weight	16.60%

97.1% patients used to take vegetables every day. 39.9% patients took red meat once in a week either beefs or mutton. History of radiation exposure (X-ray) before diagnosis of cancer was 68.7% patients, majority 48.9% had history of one X-ray, 19.8% had history of 2 X-ray and 5.8% patients had history of 5 X-ray. 80.2% of the patient had history of sun exposure, among them 34.2% exposed in sun, up to 2 h/day, 20.1% was exposed 3 h/day and 5.8% exposed 5 h/day (Table 3).

Among the study people majority were B+ blood group 43.1%, followed by A+ 25.2% and O+ blood group 17.3%. AB+, B- and AB- were in only 2.9%.

In our study, metastasis occurred at the time of diagnosis in 48.6% of the cases. 91.7% of the patients were suggested chemotherapy and the rest were both chemotherapy and radiotherapy.

Discussion

Cancer has been commonly noticed to develop in older people; and interestingly 78% of all cancer diagnoses are being conducted in people 55 years of age or older [8]. Two hospital-based (The NICRH and BSMMU) cancer registries in Bangladesh showed that 66% of the cancer patients are in the age group 30 to 65 years, the main workforce structure of a country [3]. Our study also found similar result, mean age of the cancer patient was 48.92 years. In our study male were more than female, similar findings were observed in a study done in Aizwal [9]. It has been seen that illiteracy has direct relation with the prevalence of certain cancers. Even in this study more than 54.6% patients were illiterate, followed by those who had only primary schooling. This could be ascribed to the fact that awareness pertaining to risk factors is less in illiterate people hence, were succumbing to cancer. The results of study done by Swaminathan et al. [10] also corroborated that men and women who were illiterate had a higher overall cancer incidence rates compared to the educated population. In 85.9% of the patients monthly income was less than 5000 BDT (59.52 USD), cancer management is costly in regards to drugs and radiotherapy, besides in our country Government's support is inadequate [3], so it's very difficult to complete the chemotherapy and/or radiotherapy regimen for most of the patients.

According to cancer awareness foundation [11], top five cancer of our country (according to incidence) are esophageal (13.9%), oral (8.9%), breast (8.5%), lungs (8.2%) and cervix & uterus (5.4%); our study findings were contradictory to their findings. Among all cancer, carcinoma of breast was the highest 17.89% followed by carcinoma of ovary 12.14%, but as a group GIT malignancy which includes oral cavity, tongue, esophagus, stomach, and colon were maximum 31.94%. Tobacco consumption remains the most important avoidable cancer risk. Between 25% and 30% of all cancers in developed countries is tobacco-related [12]. Smoking is indeed the most important risk factor for lung cancer, throat cancer, urinary bladder cancer, myeloid leukemia, and pancreatic cancer too [8,13-16]. Another study of Trivandrum, India emphasized that smoking increased the risk of oral cancer in men by as much as 90% [17]. In our study smoking in males was more as compared to that in females. Similar findings were evident in other research done by Murthy et al. [18]; they also showed that the major risk factors for cancer were tobacco, alcohol consumption, infections, dietary habits and behavioral risk factors. In our study 8.4% patient used to chew tobacco leaf in the form of jorda and gul, this may be one of the causes of higher incidence of oral cancer in this region. The NICRH and BSMMU cancer registry data revealed that 60% of the cancers in the male and 5% of the cancers in female are tobacco related and hence entirely avoidable [19]. So, tobacco control may be one of the most important steps to prevent cancer. Though exposure to ionizing radiation is known to increase the risk of most types of leukemia [8] in our study radiation exposure (X-ray) before diagnosis of cancer was found in 68.7% patients in the form of diagnostic X-ray.

Two hospital based study of our country revealed that disease

with distant metastasis at the time of diagnosis is <15% [3]. But in our study distant metastasis was 48.6% of the cases. This may be due to the ignorance of the people, delayed presentation and/or poor medical facilities to diagnose cancer at the vicinity. Improve cancer awareness of the people and improving medical facilities may diagnose the cancer at its earliest form.

Conclusion

Cancer is more common in older age, illiterate, poor socio-economic group and patient from rural area. Carcinoma of breast was the highest among all cancer. But as a group majority of the cases were GIT cancer. Approximately half of the cases were metastasized at the time of diagnosis.

Acknowledgement

The authors of this study are grateful to Department of Oncology of Rangpur Medical College Hospital.

References

1. Bleyer A, Barr R, Hayes-Lattin B, Thomas D, Ellis C, Anderson B, et al. The distinctive biology of cancer in adolescents and young adults. *Nat Rev Cancer*. 2008;8(4):288-98.
2. Jemal A, Center MM, DeSantis C, Ward EM. Global patterns of cancer incidence and mortality rates and trends. *Cancer Epidemiol Biomarkers Prev*. 2010;19(8):1893-907.
3. National Cancer Control Strategy and Plan of Action 2009-15. Directorate General of Health Services. Ministry of Health and Family Welfare, Bangladesh.
4. National Institute of Cancer Research and Hospital Bangladesh. 2014. National Institute of Cancer Research and Hospital Bangladesh Report 2008-2010. NICRH, Dhaka, Bangladesh.
5. Soliven K, Wang X, Small CT, Feeroz MM, Lee EG, Craig KL, et al. Simian foamy virus infection of rhesus macaques in Bangladesh: Relationship of latent proviruses and transcriptionally active viruses. *J Virol*. 2013;87(24):13628-39.
6. Hossain MA, Raton KA, Noor R. Bacteriological study of stool samples collected from children suffering from diarrhea. *J Global Biosci*. 2013;2(5):160-5.
7. Yeole BB. Role of the cancer registries in determining cancer mortality in Asia? *Asian Pac J Cancer Prev*. 2006;7(3):489-91.
8. American Cancer Society. Cancer facts and figures. *Lancet*. 2014;371(9625):1695-709.
9. Phukan RK, Zomawia E, Narain K, Hazarika NC, Mahanta J. Tobacco use and stomach cancer in Mizoram, India. *Cancer Epidemiol Biomarkers Prev*. 2005;14:1892-6.
10. Swaminathan R, Selvakumaran R, Esmay PO, Sampath P, Ferlay J, Jissa V, et al. Cancer pattern and survival in a rural district in South India. *Cancer Epidemiol*. 2009;33(5):325-31.
11. Common cancers in Bangladesh.
12. Hill C. Tobacco epidemiology. *Rev Prat*. 2012;62:325, 327-9.
13. Garavello W, Bertuccio P, Levi F, Lucchini F, Bosetti C, Malvezzi M, et al. The oral cancer epidemic in central and eastern Europe. *Int J Cancer*. 2010;127(1):160-71.
14. Wen CP, Tsai MK, Chung WS, Hsu HL, Chang YC, Chan HT, et al. Cancer risks from betel quid chewing beyond oral cancer: A multiple-site carcinogen when acting with smoking. *Cancer Causes Control*. 2010;21:1427-35.
15. D'Souza G, Agrawal Y, Halpern J, Bodison S, Gillison ML. Oral sexual

- behaviors associated with prevalent oral human papillomavirus infection. *J Infect Dis.* 2009;199(1):1263-9.
16. Jayalekshmi PA, Gangadharan P, Akiba S, Nair RR, Tsuji M, Rajan B. Tobacco chewing and female oral cavity cancer risk in Karunagappally cohort, India. *Br J Cancer.* 2009;100(5):848-52.
17. Muwonge R, Ramadas K, Sankila R, Thara S, Thomas G, Vinoda J, et al. Role of tobacco smoking, chewing and alcohol drinking in the risk of oral cancer in Trivandrum, India: A nested case-control design using incident cancer cases. *Oral Oncol.* 2008;44(5):446-54.
18. Murthy NS, Rajaram D, Gautham MS, Shivaraj NS, Nandakumar BS, Pruthvish S. Risk of cancer development in India. *Asian Pac J Cancer Prev.* 2011;12(2):387-91.
19. Cancer Registry Report National Institute of Cancer Research and Hospital 2005-07.