Clinics in Oncology

9

A Case Report of Pubic Bone Metastasis from Bladder Cancer Mimicking Osteitis Pubis

Ya-Wen Chuang¹, Chien-Chin Hsu², Chin-Chuan Chang¹, Chia-Yang Lin¹, Ying-Fong Huang^{1,3} and Yu Chang Tyan^{3,4,5,6,7*}

¹Department of Nuclear Medicine, Kaohsiung Medical University Hospital, Taiwan

²Department of Nuclear Medicine, Kaohsiung Chang Gung Memorial Hospital, Chang Gung University College of Medicine, Taiwan

³Department of Medical Imaging and Radiological Sciences, College of Health Science, Kaohsiung Medical University, Taiwan

⁴Department of Infectious Disease and Cancer Research, Kaohsiung Medical University, Taiwan

⁵Department of Medical Science and Technology, National Sun Yat-sen University, Taiwan

⁶Department of Medicine, College of Medicine, Kaohsiung Medical University, Kaohsiung, Taiwan

⁷Department of Medical Research, Kaohsiung Medical University Hospital, Kaohsiung, Taiwan

Abstract

Most symmetric radiotracer accumulation in the skeleton found in bone scintigraphy is thought to be non-specific or benign cause. Solitary symmetric bone metastasis is rare and often misinterpreted. Other image modalities or biopsy are usually needed to differentiate malignant metastasis from benign cause. The authors report a case of bladder cancer with pubic bone metastasis. The bone scan showed symmetric uptake in bilateral pubic bones mimicking the typical appearance of osteitis pubis. However, the pelvic CT scan showed bone metastasis to symphysis pubis.

Keywords: Bone scintigraphy; Bone metastasis; Bladder cancer; Osteitis pubis

Introduction

OPEN ACCESS

*Correspondence:

Yu Chang Tyan, Department of Infectious Disease and Cancer Research, Kaohsiung Medical University, Kaohsiung, Taiwan, E-mail: yctyan @kmu.edu.tw Received Date: 16 May 2017 Accepted Date: 26 Jul 2017 Published Date: 31 Jul 2017

Citation:

Chuang Y-W, Hsu C-C, Chang C-C, Lin C-Y, Huang Y-F, Tyan YC. A Case Report of Pubic Bone Metastasis from Bladder Cancer Mimicking Osteitis Pubis. Clin Oncol. 2017; 2: 1319.

Copyright © 2017 Yu Chang Tyan. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited. Osteitis pubis (also known as the pubic symphysitis) is a painful, inflammatory, noninfectious condition that involves the periosteum, bonecartilage, and ligament us structures in the symphysis pubis. It can result from several causes such as genitourinary or gynaecological surgery in older individuals [1]. One of these is mechanical strain, either from a single major event or from repetitive micro trauma, such as from running [2-7].

Case Report

A 61-year-old female patient had end stage renal disease and received hemodialysis for 5 years.



Figure 1: The symmetrically increased radioactivity insymphysis pubis and adjacent superior pubic rami, mimicking the typical appearance of symphysitis pubis.



Figure 2: The pelvic CT scan showed bone destruction of the symphysis pubis withadjacent soft tissue lesion.

Due to painless hematuria, she came to our hospital for help and transitional cell carcinoma was diagnosed. Therefore, she underwent bilateral nephrourectomy, radical cystectomy and pelvic lymph node dissection as a treatment [8].

Discussion

One year later, Tc-99mMDP bone scanand pelvic CT scan were arranged for her follow-up survey. The whole body bonescan was performed at 3 hours after the injection of 740MBq (20 mCi) Tc-99m methylene diphosphonate (MDP). The symmetrically increased radioactivity in symphysis pubis and adjacent superior pubic rami, mimicking the typical appearance of symphysitis pubis (Figure 1).

The pelvic CT scan showed bone destruction of the symphysis pubis withadjacent soft tissue lesion (Figure 2). Metastatic disease is considered. In summary, this case illustrates the potential pitfalls of misinterpreting solitary symmetric bone metastasis as benign cause. Careful attention to the clinical and radiographic presentation is important to avoid this misdiagnosis.

Acknowledgments

This work was supported by Research Grants MOST 103-2320-B-037-025 from the Ministry of Science and Technology, KMU-TP105E12, KMU-TP105PR06, KMU-M1060289 from Kaohsiung Medical University, and NSYSUKMU106-P011 from NSYSU-KMU Research Project, Taiwan.

References

- Gleichmann R, Gasirabo P. Perineal endoscopic desobliteration of a complete anastomotic stricture following a salvage radical prostatectomy. Aktuelle Urol. 2011; 42(6): 55-57.
- Major NM, Helms CA. Pelvic stress injuries: the relationship between osteitis pubis (symphysis pubis stress injury) and sacroiliac abnormalities in athletes. Skeletal Radiol. 1997; 26(2):711-717.
- Kanstrup IL. Bone scintigraphy in sports medicine: a review. Scand J Med Sci Sports. 1997; 7(2): 322-330.
- Koulouris G. Imaging review of groin pain in elite athletes: an anatomic approach to imaging findings. AJR Am J Roentgenol. 2008; 191(4): 962-972.
- Rex C, Elsworth C. Pelvic insufficiency fracture with diastasis of the pubic symphysis after irradiation: a case report. J Bone Joint Surg Br. 1998; 80(2): 264-266.
- Ikushima H, Osaki K, Furutani S. Pelvic bone complications following radiation therapy of gynecologic malignancies: clinical evaluation of radiation-induced pelvic insufficiency fractures. Gynecol Oncol. 2006; 103(2): 1100-1104.
- 7. Koch RA, Jackson DW. Pubic symphysitis in runners: a report of two cases. Am J Sports Med. 1981; 9(2): 62-63.
- Wu CF, Shee JJ, Ho DR. Different treatment strategies for end stage renal disease in patients with transitional cell carcinoma. J Urol. 2004; 171(2): 126-129.